

PHARMACOR PROZINE 5

Prochlorperazine maleate

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Pharmacor Prozine 5.

It does not contain all the available information.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking Pharmacor Prozine 5 against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine. You may need to read it again.

What Pharmacor Prozine 5 is used for

Pharmacor Prozine 5 belongs to a group of medicines called phenothiazines. It helps to correct chemical imbalances in the brain, allowing it to function correctly. These chemicals may also affect the parts of the brain which control nausea (feeling sick) and vomiting.

Pharmacor Prozine 5 is used to treat nausea, vomiting and dizziness due to various causes, including migraine (severe headache).

Your doctor may have prescribed Pharmacor Prozine 5 for another reason.

Ask your doctor if you have any questions about why Pharmacor Prozine 5 has been prescribed for you.

This medicine is available only with a doctor's prescription.

Pharmacor Prozine 5 is not recommended for use in children (under the age of 2 years or children under 10 kg in weight)

Before you take Pharmacor Prozine 5

When you must not take it

Do not take Pharmacor Prozine 5 if you have an allergy to:

- Prochlorperazine
- The group of medicines called phenothiazines
- Any of the ingredients listed at the end of this leaflet

Symptoms of an allergic reaction to Pharmacor Prozine 5 may include:

- Shortness of breath, wheezing or difficulty in breathing
- Swelling of the face, lips, tongue or other parts of the body
- Rash, itching or hives on the skin

You should not take Pharmacor Prozine 5 if you have any of the following medical conditions:

- Shock
- Disease of the blood with a low number of blood cells
- Yellowing of the skin and/or eye, also called jaundice

Pharmacor Prozine 5 must not be given to anyone who is unconscious or in a coma.

Do not take any medicines that cause drowsiness while you are taking Pharmacor Prozine 5.

Do not take Pharmacor Prozine 5 after the expiry date (EXP) printed on the pack.

If you take this medicine after the expiry date has passed, it may not work as well.

Do not take Pharmacor Prozine 5 if the packaging is torn or shows signs of tampering.

If you are not sure whether you should start taking Pharmacor Prozine 5, talk to your doctor or pharmacist.

Before you start to take it

Tell your doctor or pharmacist if you have allergies to:

- Any other medicines
- Any other substances, such as foods, preservatives or dyes.

Tell your doctor if you are pregnant or intend to become pregnant.

Like most phenothiazine medicines, Pharmacor Prozine 5 is not recommended for use during pregnancy. If there is a need to take Pharmacor Prozine 5 during your pregnancy, your doctor will discuss with you the benefits and risks of using it.

Tell your doctor or pharmacist if you are breastfeeding or plan to breastfeed.

It is recommended that you do **not** breastfeed while taking Pharmacor Prozine 5, as it is not known whether Pharmacor Prozine 5 passes into breast milk.

Tell your doctor or pharmacist if you have or have had any medical conditions, especially the following:

- Phaechromocytoma, a rare tumour of the adrenal glands which sit near the kidneys.
- Parkinson's disease, a disease of the brain affecting movement which causes trembling, rigid posture, slow movement and a shuffling, unbalanced walk.
- Myasthenia gravis, a disease of the muscles causing drooping eyelids, double vision, difficulty in speaking and swallowing and sometimes muscle weakness in the arms or legs.
- Kidney problems
- Heart and blood vessel problems, low blood pressure, blood clots
- Liver disease
- Prostate problems
- Epilepsy, seizures or fits
- Low blood calcium levels
- An overactive thyroid gland
- Glaucoma, a condition in which there is a build-up of fluid in the eye.
- Neuroleptic Malignant Syndrome, a reaction to some medicines with a sudden increase in body temperature, extremely high blood pressure and severe convulsions.
- A reaction to some medicines with uncontrollable twitching or jerking movements of the arms and legs.
- Dementia
- High blood sugar levels

Tell your doctor if you are about to have any surgery which requires a general anaesthetic.

If you have not told your doctor or pharmacist about any of the above, tell them before you start taking Pharmacor Prozine 5.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Pharmacor Prozine 5 may interfere with each other. These include:

- Some medicines used to control depression or mood swings
- Alcohol
- Desferrioxamine, a drug used in iron overdose
- Procarbazine, an anticancer drug
- Some medicines used to control epilepsy
- Medicines used to treat Parkinson's disease
- Anticholinergic medicines which are used to relieve stomach cramps, spasms and travel sickness
- Atropine, a medicine which may be used in some eye drops or cough and cold preparations
- Some oral medicines used to prevent your blood from clotting
- Medicines used to treat high blood pressure and fluid build-up in your body
- Other medications such as bepridil, cisapride, sultopride, thioridazine, methadone, erythromycin injection, vincamine injection, halofantane, pentamidine, sparfloxacin, amphotericin B, glucocorticoids or tetracosactides
- Anitpsychotics
- Certain laxatives

These medicines may be affected by Pharmacor Prozine 5 or may affect how well it works. You may need different amounts of your medicine or you may need to take different medicines. Your doctor will advise you.

Your doctor and pharmacist may have more information on medicines to be careful with or avoid while taking Pharmacor Prozine 5.

How to take Pharmacor Prozine 5

Follow all directions given to you by your doctor and pharmacist carefully.

They may differ from the information contained in this leaflet.

If you do not understand the instructions on the pack, ask your doctor or pharmacist for help.

How much to take

Your doctor or pharmacist will tell you how much Pharmacor Prozine 5 you will need to take each day. This depends on your condition and whether or not you are taking any other medicines.

The usual recommended dose for nausea and vomiting is 1 or 2 tablets two to three times daily.

The usual recommended dose for dizziness is 1 or 2 tablets three to four times daily.

How to take it

Swallow Pharmacor Prozine 5 tablets whole with a glass of water. Do not chew the tablets.

When to take it

It does not matter if you take Pharmacor Prozine 5 before or after food.

If you are not sure what to do, ask your doctor or pharmacist.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

How long to take it

Continue taking Pharmacor Prozine 5 for as long as your doctor tells you.

If you take too much (overdose)

Do not try to vomit.

Immediately telephone your doctor or pharmacist or the Poisons Information Centre (13 11 26) or go to Accident and Emergency at your nearest hospital if you think that you, or anyone else, has taken too much Pharmacor Prozine 5. Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

Symptoms of overdose may include the following:

- Coma
- Restlessness, shaking, muscle twitching, muscle weakness, spasm
- Confusion
- Excitement or agitation
- Low blood pressure
- Fast heart beat
- Decrease in body temperature
- Small pupils in the eye
- Difficulty in swallowing or breathing
- Blue skin

Your doctor or pharmacist has information on how to recognize and treat an overdose. Ask your doctor or pharmacist if you have any concerns.

While you are taking Pharmacor Prozine 5

Things you must do

Tell your doctor immediately if you notice any uncontrolled movements of the tongue, face, mouth or jaw, such as puffing of the cheeks, puckering of the mouth or chewing movements.

These are symptoms of a very rare condition called Tardive Dyskinesia, which may develop in people taking phenothiazine medicine, including Pharmacor Prozine 5.

The condition is more likely to occur during long term treatment with Pharmacor Prozine 5, especially in elderly women. In very rare cases, this may be permanent.

Tell any other doctors, dentists and pharmacists who are treating you that you are taking Pharmacor Prozine 5.

If you are about to be started on any new medicines, tell your doctor, dentist or pharmacist that you are taking Pharmacor Prozine 5.

If you plan to have surgery that needs a general anaesthetic, tell your doctor or dentist that you are taking Pharmacor Prozine 5.

If you become pregnant while taking Pharmacor Prozine 5, tell your doctor immediately.

Things you must not do

Do not give Pharmacor Prozine 5 to anyone else, even if they have the same condition as you.

Do not take Pharmacor Prozine 5 to treat any other complaints unless your doctor or pharmacist tells you to.

Do not stop taking Pharmacor Prozine 5, or lower the dosage, even if you are feeling better, without checking with your doctor.

If you stop taking Pharmacor Prozine 5 suddenly, your condition may worsen or your chance of getting an unwanted side effect may increase. To prevent this, your doctor may gradually reduce the amount of Pharmacor Prozine 5 you take each day before stopping completely.

Things to be careful of

Be careful driving or operating machinery until you know how Pharmacor Prozine 5 affects you.

As with other medicines, Pharmacor Prozine 5 may cause dizziness, light-headedness, tiredness, drowsiness in some people. Make sure you know how you react to Pharmacor Prozine 5 before you drive a car, operate machinery, or do anything else that could be dangerous if you are dizzy or light-headed. If this occurs do not drive. If you drink alcohol, dizziness or light-headedness may be worse.

If Pharmacor Prozine 5 makes you feel light-headed, dizzy or faint, be careful when getting up from a sitting or lying position. Getting up slowly may help.

Be careful when drinking alcohol while taking Pharmacor Prozine 5. Combining Pharmacor Prozine 5 and alcohol can make you more sleepy, dizzy or light-headed.

Your doctor may suggest you avoid alcohol while you are being treated with Pharmacor Prozine 5.

If outdoors, wear protective clothing and use at least a 15+ sunscreen.

Pharmacor Prozine 5 may cause your skin to be much more sensitive to sunlight than it is normally. Exposure to sunlight may cause a skin rash, itching, redness, or even severe sunburn. If your skin does appear to be burning, tell your doctor.

Make sure you keep cool in hot weather and keep warm in cool weather.

Pharmacor Prozine 5 may affect the way your body reacts to temperature changes.

Side Effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Pharmacor Prozine 5.

Pharmacor Prozine 5 helps most people with nausea, vomiting and dizziness, but it may have unwanted side effects in a few people.

All medicines can have side effects.

Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects. If you are over 65 years of age you may have an increased chance of getting side effects.

Ask your doctor or pharmacist to answer any questions you may have.

If you get any side effects, do not stop taking Pharmacor Prozine 5 without first talking to your doctor or pharmacist.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- Constipation
- Dry mouth
- Drowsiness
- Restlessness
- Trembling, rigid posture, mask-like face, slow movements and a shuffling unbalanced walk
- Twitching
- Blurred vision

The following side effects are less common:

- Low blood pressure
- Changes in heart beats
- Swelling of the hands, ankles or feet
- Skin rash

- For females: unusual secretion of breast milk, irregular periods
- For males: breast enlargement, difficulty in ejaculating
- Severe pain in the stomach with bloating, cramps and vomiting
- Difficulty passing urine
- Yellowing of the skin and/or eyes
- Headache
- Insomnia
- Seizures
- Difficulty in breathing
- High blood sugar levels

If any of the following happen, tell your doctor or pharmacist immediately or go to Accident and Emergency at your nearest hospital:

- Unusual muscle tone or spasms causing distortion of the body in children

These are very serious side effects.

You may need urgent medical attention or hospitalization.

All of these side effects are very rare.

Other side effects not listed above may occur in some patients. Tell your doctor or pharmacist if you notice anything that is making you feel unwell.

Do not be alarmed by this list of possible side effects.

You may not experience any of them.

After using Pharmacor Prozine 5

Storage

Keep Pharmacor Prozine 5 tablets in a cool dry place where the temperature stays below 25°C. Protect from light.

Keep your Pharmacor Prozine 5 in the pack until it is time to take them.

If you take the tablets out of the pack, they may not keep well.

Do not store Pharmacor Prozine 5 or any other medicine in the bathroom or near a sink.

Do not leave it in the car on hot days or on window sills.

Heat and dampness can destroy some medicines.

Keep it where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor or pharmacist tells you to stop taking Pharmacor Prozine 5, or the medicine has passed its expiry date, ask your pharmacist what to do with any that are left over.

Product Description

What it looks like

Pharmacor Prozine 5 tablets are white, round and marked with "5". They are available in packs of 10 and 25 tablets.

Ingredients

Pharmacor Prozine 5 tablets contain: Prochlorperazine maleate 5mg (active ingredient), lactose, maize starch, purified water, colloidal anhydrous silica, magnesium stearate.

Supplier

Pharmacor Limited
Dee Why NSW 2099

Registration number:
AUST R 172127

Date of preparation:
February 2011