

# Exemestane AN

Exemestane

---

## Consumer Medicine Information

**Please read this leaflet carefully before you start Exemestane AN.**

---

### What is in this leaflet

---

This leaflet answers some common questions about Exemestane AN.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking Exemestane AN against the benefits it is expected to have for you.

**If you have any concerns about this medicine, ask your doctor or pharmacist.**

**Keep this leaflet.**

You may need to read it again.

---

### What Exemestane AN is used for

---

Exemestane AN is used to treat breast cancer in women who no longer have their menstrual periods, either naturally due to their age (after menopause) or because they have had their ovaries surgically removed.

Exemestane AN is used to reduce the risk of recurrence or spreading of the breast cancer. It is also used when the cancer has not responded or has returned after treatment with another medicine or medicines.

### *How Exemestane AN works*

Exemestane AN is an aromatase inactivator. It works by significantly reducing the supply of oestrogen to cancer cells.

This stops the growth of those cancer cells which need oestrogen, a natural female sex hormone, to grow.

**Ask your doctor if you have any questions about why Exemestane AN has been prescribed for you.**

Your doctor may have prescribed it for another purpose.

Exemestane AN is only available with a doctor's prescription. It is not addictive.

---

### Before you take it

---

#### *When you must not take it*

**Do not take Exemestane AN and talk to your doctor if you think you may have an allergy to:**

- exemestane, the active ingredient in Exemestane AN
- any of the other ingredients in Exemestane AN listed at the end of this leaflet.

Symptoms of an allergic reaction may include shortness of breath, wheezing or difficulty breathing; swelling of the face, lips, tongue or other parts of the body; rash, itching or hives on the skin.

**Do not take Exemestane AN if you are pregnant or intend to become pregnant.**

Exemestane AN may affect your developing baby if you take it during pregnancy.

**Do not breastfeed while taking Exemestane AN.**

There is a possibility that your baby may be affected.

Exemestane AN is not recommended for use in children.

**Do not take Exemestane AN after the expiry date printed on the pack.**

**Do not take Exemestane AN if the packaging is torn or shows signs of tampering.**

**If your Exemestane AN has expired or is damaged return it to your pharmacist for disposal.**

**If you are not sure whether you should start taking Exemestane AN, talk to your doctor.**

#### *Before you start to take it*

**Tell your doctor if you have an allergy to:**

- any medicine
- other substances, such as foods, preservatives or dyes.

**Tell your doctor if you:**

- are still having your monthly period
- have or have had kidney problems
- have or have had liver problems
- have osteoporosis (disease which causes bones to be more brittle and likely to break)

**If you have not told your doctor about any of the above, tell your doctor before you start taking Exemestane AN.**

## *Taking other medicines*

**Tell your doctor or pharmacist if you are taking any other medicines, including medicines that you buy without a prescription from a pharmacy, supermarket or health food shop.**

Some medicines should not be used with Exemestane AN. These include:

- medicines which contain oestrogen, such as hormone replacement therapy (HRT)
- oral contraceptives

Some health food products for menopausal symptoms contain natural oestrogens.

**Tell your doctor or pharmacist if you are taking any medicines or health food products containing oestrogens.**

**Talk to your doctor or pharmacist if you have any concerns or questions about taking Exemestane AN.**

---

## How to take it

---

**Follow all directions given to you by your doctor or pharmacist carefully.**

They may differ from the information contained in this leaflet.

**If you do not understand the instructions on the box, ask your doctor or pharmacist for help.**

### *How much to take*

The recommended dose is one 25 mg tablet taken once daily.

### *When to take it*

**Take the tablet once daily after a meal at approximately the same time each day.**

### *How long to take it*

Your doctor or pharmacist will tell you how long to take Exemestane AN.

**Do not stop taking Exemestane AN unless your doctor tells you to, even if you feel better.**

### *If you forget to take it*

**If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to. Otherwise, take it as soon as you remember, and then go back to taking Exemestane AN as you would normally.**

**Do not take a double dose to make up for the one that you missed.**

**If you have trouble remembering to take your medicine, ask your pharmacist for some hints.**

### *If you take too much (overdose)*

**Immediately telephone your doctor or Poisons Information Centre (Phone Australia 13 11 26) for advice or go to Accident and Emergency at your nearest hospital, if you think you or anyone else may have taken too much Exemestane AN.**

**Do this even if there are no signs of discomfort or poisoning.**

**Keep the telephone numbers for these places handy.**

---

## While you are taking it

---

### *Things you must do*

**Use Exemestane AN exactly as your doctor has prescribed.**

**Be sure to keep all your doctor's appointments so your progress can be checked.**

**If you feel that your medicine is not helping your condition, talk to your doctor.**

**Tell all doctors, dentists and pharmacists who are treating you that you are taking Exemestane AN.**

**If you go into hospital whilst taking Exemestane AN, let the medical staff know that you are taking Exemestane AN.**

**If you are about to be started on any new medicine, tell your doctor, dentist or pharmacist that you are taking Exemestane AN.**

### *Things you must not do*

**Do not give Exemestane AN to anyone else, even if they have the same condition as you.**

**Do not take Exemestane AN to treat any other complaints unless your doctor tells you to.**

**Do not stop taking Exemestane AN or change the dosage without checking with your doctor.**

### *Things to be careful of*

**Be careful driving or operating machinery, until you know how Exemestane AN affects you.**

Exemestane AN may cause tiredness or dizziness in some people.

**If you experience these effects whilst taking Exemestane AN, you should not attempt to drive or operate machinery.**

**If Exemestane AN makes you feel dizzy, be careful when getting up from a sitting or lying position.**

---

## Side effects

---

**Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Exemestane AN.**

Exemestane AN, like other medicines, may cause unwanted side effects in some people. Many women can take Exemestane AN without any problems, but some women may have mild to moderate side effects.

**Tell your doctor or pharmacist as soon as possible if you have any of the following side effects or other effects that you think may be related to your treatment:**

- hot flushes

- nausea, vomiting, loss of appetite, indigestion, pain (including stomach ache)
- constipation
- diarrhoea
- tiredness
- swelling in your legs
- dizziness and headache
- increased sweating
- difficulty sleeping
- depression
- disturbed vision such as blurriness
- pain in your muscle or joints
- increase in weight
- skin rash
- hair loss
- bleeding or bruising more easily than normal
- pain and/or numbness of hands, loss of feeling in fingers/thumb
- frequent infections such as fever, severe chills, sore throat or mouth ulcers
- symptoms of gastric ulcer such as blood in stools, black tarry stools or vomiting of blood
- abnormal vaginal bleeding
- hepatitis, yellowing of the skin or eyes, also called jaundice.

Other side effects not listed above may occur in some people. Some of these side effects can only be found when your doctor does tests from time to time to check your progress.

**Do not be alarmed by this list of possible side effects.**

You may not experience any of them.

---

## After using it

---

### *Storage*

**Keep your tablets in their blister pack until it is time to take them.**

If you take the tablets out of the blister pack they may not keep well.

**Keep Exemestane AN in a cool dry place where the temperature stays below 30°C.**

**Do not leave Exemestane AN or any other medicine in the car or on window sills.**

**Do not store Exemestane AN or any other medicines in a bathroom or near a sink.**

Heat and dampness can destroy some medicines.

**Keep Exemestane AN where children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

### *Disposal*

**If your doctor tells you to stop taking Exemestane AN or the tablets have passed their expiry date, ask your pharmacist what to do with any tablets that are left over.**

---

## Product description

---

### *What it looks like*

Exemestane AN 25 mg tablets are white to off-white, circular, biconvex and are sugar-coated.

### *Ingredients*

#### **Active Ingredient**

Exemestane AN tablets contain 25 mg of exemestane as the active ingredient.

#### **Other Ingredients**

- Microcrystalline cellulose,
- crospovidone,
- polysorbate 80,
- mannitol,
- anhydrous colloidal silica,
- magnesium stearate,
- sucrose,
- acacia,
- purified talc,
- titanium dioxide,
- shellac,
- beeswax white, and
- carnauba wax yellow.

### *Registration Number*

AUST R 177311

### *Sponsor*

**Southern Cross Pharma Pty Ltd**  
56 Illabunda Drive  
Malua Bay NSW 2536  
ABN 47 094 447 677

### *Date of preparation*

February 2014