

SOLIRIS®

Eculizumab (rmc)

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about SOLIRIS. It does not contain all the available information.

It does not take the place of talking to your doctor, nurse or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you having SOLIRIS against the benefits they expect it will have for you.

If you have any concerns about this medicine, ask your doctor or nurse.

Keep this leaflet.

You may need to read it again.

What SOLIRIS is used for

SOLIRIS is a medicine containing an active substance called eculizumab (rmc) which belongs to a class of medicines called monoclonal antibodies.

SOLIRIS is used for the treatment of patients with a disease that affects red blood cells called Paroxysmal Nocturnal Haemoglobinuria (PNH).

SOLIRIS is also used to treat patients with a condition called atypical Haemolytic Uraemic Syndrome (aHUS).

How it works

Patients with PNH lack naturally occurring protective proteins on the surface of some of their red blood cells. In unaffected individuals, these proteins protect red blood

cells from destruction by the body's inflammatory response. PNH patients lack these protective proteins and their red blood cells can be destroyed. This can lead to low blood counts (anaemia), tiredness, difficulty in functioning, pain, dark urine, kidney failure, shortness of breath, and blood clots.

SOLIRIS can block the body's inflammatory response, and its ability to attack and destroy its own vulnerable PNH blood cells. In this way SOLIRIS improves anaemia, fatigue, and other signs and symptoms of PNH.

Patients with aHUS have an inflammatory condition which affects the blood system and kidney. This can lead to reduced or lost kidney function, blood clots, low blood counts (low platelets and anaemia), tiredness and difficulty functioning. SOLIRIS works by blocking the body's inflammatory response and its ability to attack and destroy its own vulnerable blood and kidney cells.

Before you are given SOLIRIS

When you must not be given SOLIRIS

SOLIRIS treatment may reduce your natural resistance to infections, especially against certain organisms that cause meningitis (infection of the linings of the brain).

DO NOT use SOLIRIS if:

- you have not been vaccinated against meningococcal infections,
- you are not up to date with your meningococcal vaccination,
- if it is less than 2 weeks after receiving your meningococcal vaccination and you are not taking antibiotics to reduce the risk of infection,
- you have meningitis

Do not use SOLIRIS if you have had an allergic reaction to:

- SOLIRIS, or any of the ingredients listed at the end of this leaflet, or
- any other proteins of mouse origin

Symptoms of an allergic reaction may include;

- shortness of breath, wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin

If you are not sure whether you should be treated with SOLIRIS, talk to your doctor or nurse.

Before you start SOLIRIS treatment

You must be aware of the following signs and symptoms of a meningococcal infection:

- headache with nausea or vomiting
- headache with a stiff neck or stiff back
- fever
- rash
- confusion
- severe muscle aches with flu-like symptoms

- sensitivity to light

Call your doctor immediately and go to Accident and Emergency at your nearest hospital if you have any of the symptoms listed above.

Patient Safety Card

Because of the importance of rapidly identifying and treating certain types of infection you will be provided with a Patient Safety Card.

You must carry this card with you at all times and show it to any doctor or nurse that treats you.

You must receive a meningococcal vaccine before or at the time of your first dose of SOLIRIS unless you have already been vaccinated with this vaccine or you take antibiotics to reduce the risk of infection until 2 weeks after you have been vaccinated.

If you have been vaccinated with a meningococcal vaccine in the past, you might need a booster dose before or at the time of starting SOLIRIS.

Your doctor will decide if you need another dose of a meningococcal vaccine.

You should also be aware that vaccination may not prevent this type of infection.

You may need antibiotics to prevent infection.

If you are less than 18 years old, you must also be vaccinated against Haemophilus influenzae and pneumococcal infections.

Your doctor will arrange this according to national vaccination recommendations for your age group.

Tell your doctor if you have an infection.

SOLIRIS may reduce your natural resistance to infection.

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Allergic or anaphylactic (more severe allergic) reactions may occur with SOLIRIS treatment (known as infusion related reactions). Your

doctor or nurse will check for side effects during your infusion and for one hour afterwards. See "Side Effects" for symptoms to look out for.

Tell your doctor if you are pregnant or want to become pregnant.

SOLIRIS has not been studied in pregnant women. Women who are able to get pregnant should use effective contraception during treatment, and up to 5 months after treatment.

Tell your doctor if you are breastfeeding.

It is not known whether SOLIRIS passes into breast milk. Since many medicines are secreted into breast milk, breastfeeding should be discontinued during treatment, and up to 5 months after treatment.

Tell your doctor if you are on a sodium controlled diet.

SOLIRIS contains 115 mg sodium per vial, which may need to be considered in calculating your sodium intake.

If you have not told your doctor or nurse about any of the above, tell them before you are given SOLIRIS.

Taking other medicines

Tell your doctor or nurse if you are taking any other medicines, including any that you buy without prescription from your pharmacy, supermarket or health food shop.

The effect of using SOLIRIS on other medicines has not been studied. Ask your doctor or nurse if you have any questions.

How SOLIRIS is given

Follow all directions given to you by your doctor or nurse carefully.

They may differ from the information contained in this leaflet.

SOLIRIS will be given to you directly into the vein

(intravenously) by a doctor or nurse. Each infusion will take approximately 25 - 45 minutes.

- For adults, SOLIRIS is given once a week for the first four weeks, on the fifth week, the dose is increased and then SOLIRIS is given every two weeks
- For children and adolescents who weigh less than 40 kg, SOLIRIS will be given at a frequency and dose that varies depending on their weight
- Children and adolescents who weigh more than 40 kg are treated with the adult dosing
- If you are having plasma exchange or plasma infusion you may receive additional doses of SOLIRIS

If you miss a dose

If you forget or miss your appointment for a SOLIRIS infusion, contact your doctor immediately.

If you are given too much (overdose)

There have been no reported overdoses of SOLIRIS. As SOLIRIS is given to you under the supervision of your doctor, it is unlikely that you will receive too much.

While you are using SOLIRIS

Things you must do

Tell any other doctors, dentists, and pharmacists who treat you that you are taking this medicine.

Keep appointments with your doctor or clinic.

It is important to have the infusion with SOLIRIS at the appropriate time to make sure the medicine has the best chance of providing treatment for the condition.

Have any tests when your doctor says to.

Your doctor may wish to test your body's response to SOLIRIS to make sure that it is working or may wish to test your body's response if you stop therapy.

Things you must not do

Do not stop taking SOLIRIS without checking with your doctor.

If you forget or miss a SOLIRIS infusion, call your doctor immediately.

If you have PNH, stopping treatment with SOLIRIS may cause a sudden and serious breakdown of your red blood cells.

Symptoms or problems from red blood cell breakdown include:

- a large drop in your red blood cell count causing anaemia. Symptoms include tiredness, headaches, being short of breath when exercising, dizziness and looking pale
- confusion or change in how alert you are
- chest pain or angina
- dark urine
- blood clots

If you experience any of these symptoms, contact your doctor immediately.

Your doctor will need to monitor you closely for at least 8 weeks after stopping SOLIRIS.

If you have aHUS, stopping treatment with SOLIRIS may cause small blood clots (known as thrombotic microangiopathy or TMA).

Your doctor will discuss the possible side effects with you and explain the risks. Your doctor will want to monitor you closely.

Symptoms or problems from TMA may include:

- low blood platelet count, leading to bruising or bleeding more easily than normal

- a large drop in your red blood cell count causing anaemia. Symptoms include tiredness, headaches, being short of breath when exercising, dizziness and looking pale
- confusion or change in how alert you are
- seizures
- chest pain or angina
- decreased urination (kidney problems)
- shortness of breath
- blood clots

If you experience any of these symptoms, contact your doctor immediately.

Side Effects

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Your doctor has weighed the risks of using this medicine against the benefits they expect it will have for you.

Do not be alarmed by this list of possible side effects.

You may not experience any of them.

Tell your doctor if you notice any of the following and they worry you:

- headaches
- dizziness or lightheadedness
- runny or stuffy nose and colds
- sore throat or mouth ulcers
- cough

The above list includes the more common side effects of SOLIRIS.

Tell your doctor as soon as possible if you notice any of the following:

- bleeding or bruising more easily than normal
- tiredness or unusual weakness
- flu-like symptoms such as fever, sore throat, cough and chills

- being short of breath (especially when exercising)
- looking pale
- confusion or change in how alert you are
- difficulty breathing
- chest tightness
- chest pain or angina
- fast or irregular heartbeats, also called palpitations
- dark urine or blood in the urine
- difficulties or pain when urinating
- diarrhoea
- nausea or vomiting
- stomach pain or discomfort
- constipation
- indigestion or an uncomfortable feeling in the stomach or belching after eating
- rash or itchy skin
- hair loss
- joint pain
- back or neck pain
- muscle spasms or aching muscles, muscle tenderness or weakness not caused by exercise
- loss of appetite
- loss of taste
- pins and needles in your hands or feet
- depression, anxiety or mood swings
- difficulty sleeping, abnormal dreams

Tell your doctor or nurse immediately if you experience any side effects during or after your SOLIRIS infusion.

Infusion related reactions are not common but they may be serious. Your doctor or nurse will monitor you for one hour after your infusion.

Tell your doctor as soon as possible if you notice signs of an infection.

Examples of infection include;

- upper or lower respiratory tract infection (e.g. bronchitis or pneumonia)
- cold sores (herpes simplex)

- urinary tract infection (UTI) or cystitis
- gastro or stomach flu
- gum or tooth infections
- viral infection

SOLIRIS may increase your susceptibility to infection. Some infections are serious and can be life-threatening. You may need urgent medical attention.

Tell your doctor immediately and go to Accident and Emergency at your nearest hospital if you experience any of the following symptoms:

- headache with nausea or vomiting
- headache with a stiff neck or stiff back
- fever
- rash
- confusion
- severe muscle aches with flu-like symptoms,
- sensitivity to light

These are possible symptoms of meningococcal infection. If you have meningococcal infection you need urgent medical attention.

Always carry your SOLIRIS Patient Safety Card which lists the symptoms of meningococcal disease and important contact information.

If you get any side effects, do not stop SOLIRIS without first talking to your doctor

Ask your doctor or nurse to answer any questions you may have.

Product Description

What SOLIRIS looks like

SOLIRIS is a clear, colourless, solution contained in a 30 mL glass vial.

Ingredients

Active ingredient

- eculizumab (rmc)

Other ingredients

- sodium chloride
- monobasic sodium phosphate monohydrate
- dibasic sodium phosphate heptahydrate
- polysorbate 80
- water for injections

Manufacturer/Supplier

In Australia this product is registered by:

Alexion Pharmaceuticals
Australasia Pty Ltd
Suite 401, Level 4, Building A.
20 Rodborough Rd. Frenchs Forest.
NSW 2086.

Medical enquiries: 1800 788 189

AUST R 138885

This leaflet was revised in June 2016

Storing SOLIRIS

SOLIRIS will be stored under refrigeration (2°C to 8°C) in the hospital or pharmacy.