

TOPIRAMATE GH

Topiramate film-coated tablets

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Topiramate GH. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risk of you taking Topiramate GH against the benefits it is expected to have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

What Topiramate GH is used for

Topiramate GH is used to treat various types of epilepsy in adults and children over 2 years of age, and to prevent migraines in adults.

Topiramate GH contains the active ingredient topiramate.

Topiramate belongs to a group of medicines called anti-convulsants.

It prevents seizures and migraines by acting on the nerves and chemicals in the brain.

Your doctor may prescribe Topiramate GH on its own, or in addition to another medicine for controlling your seizures or migraines.

Ask your doctor if you have any questions about why this medicine has been prescribed for you.

Your doctor may have prescribed it for another reason.

Before you take Topiramate GH

When you must not take it

Do not take this medicine if you have an allergy to:

- the active ingredient topiramate or to any of the other ingredients listed at the end of this leaflet under Product Description;
- any other similar medicines.

Some of the symptoms of an allergic reaction may include:

- shortness of breath;
- wheezing or difficulty breathing;
- swelling of the face, lips, tongue or other parts of the body;
- rash, itching or hives on the skin.

Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should start taking this medicine, talk to your doctor.

Before you start to take it

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if you have or have ever had any of the following medical conditions:

- a personality disorder or mental illness;
- kidney stones, kidney disease or a family history of kidney stones;
- liver disease;
- eye problem or high pressure in the eye.

Tell your doctor if you are pregnant or plan to become pregnant.

Topiramate GH has caused harm to the developing foetus in animal studies. Its safety has not been verified in pregnant humans. Information available suggests that there may be an association between the use of Topiramate GH in humans during pregnancy and birth defects including cleft lip/palate.

However, it is very important to control your fits while you are pregnant. If it is necessary for you to take this medicine, your doctor can help you decide whether or not to take it during pregnancy.

Tell your doctor if you are breastfeeding or wish to breastfeed.

Topiramate GH may appear in breast milk and it is not recommended to breastfeed while taking this medicine.

If you have not told your doctor about any of the above, tell him/her before you start taking Topiramate GH.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other

medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Topiramate GH may interfere with each other. These include:

- other medicines for epilepsy, such as phenytoin or carbamazepine;
- digoxin (used to treat heart disease);
- oral contraceptives (the pill);
- metformin, pioglitazone, glibenclamide (used to treat diabetes);
- acetazolamide (used to treat glaucoma);
- lithium, risperidone (used to treat bipolar disorder and schizophrenia);
- hydrochlorothiazide (used to treat swelling or high blood pressure);
- propranolol;
- valproic acid;
- diltiazem (used to treat hypertension and prevent angina);
- venlafaxine (used to treat depression);
- any medicine which slows your reactions. This may include medicines to help you sleep or relieve pain, antidepressants, tranquillisers or antihistamines which can make you drowsy.

These medicines may be affected by Topiramate GH, or may affect how well it works. You may need different amounts of your medicines, or you may need to take different medicines.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking this medicine.

How to take Topiramate GH

Follow all directions given to you by your doctor or pharmacist carefully.

They may differ from the information contained in this leaflet.

If you do not understand the instructions, ask your doctor or pharmacist for help.

How much to take

The doses shown below are the usual recommended doses. However, your doctor may tell you to take higher or lower doses. Your doctor will start with a low dose and slowly increase the dose to the lowest amount needed to control your epilepsy.

At the start of treatment Topiramate GH may be taken once a day, preferably at night. After the dose is increased, it is taken twice a day.

You should continue taking Topiramate GH until your doctor tells you to stop. Before stopping, it is recommended that the dose be reduced gradually.

Adults

Epilepsy - Treatment starts at a low dose of 25 mg to 50 mg per day. The dose is then gradually increased by 25 mg to 100 mg over weekly periods or longer, until the most suitable dose is reached.

Migraine - Treatment starts at a low dose of 25 mg nightly for one week. The dose is then increased over weekly periods or longer by 25 mg/day, until the most suitable dose is reached.

Children (2 years and over) Epilepsy Only

Treatment starts at 25 mg or less per day, depending on the body weight. This dose is then gradually increased over weekly periods or longer, until the most suitable dose is reached.

Ask your doctor or pharmacist if you are unsure of the correct dose for you.

They will tell you exactly how much to take.

Follow the instructions they give you.

If you take the wrong dose, Topiramate GH may not work as well and your problem may not improve.

How to take it

You can take the tablets with or without food. Always swallow the tablets whole with plenty of water.

If you forget to take it

Take your dose as soon as you remember, and continue to take it as you would normally.

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Do not take a double dose to make up for the dose you missed.

This may increase the chance of you getting an unwanted side effect.

If you have missed more than one dose, or are not sure what to do, ask your doctor or pharmacist.

If you have trouble remembering when to take your medicine, ask your pharmacist for some hints.

If you take too much (overdose)

Immediately telephone your doctor or the Poisons Information Centre (telephone 13 11 26 or New Zealand 0800 POISON or 0800 764766) for advice, or go to Accident and Emergency at the nearest hospital, if you think you or anyone else may have taken too much Topiramate GH. Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

If you take too much Topiramate GH, you may experience headache, dizziness, lightheadedness, drowsiness, convulsions, speech

disturbances, double or blurred vision, difficulty with thinking, abnormal coordination, stomach pain, depression, agitation, faster breathing or you may lose consciousness.

While you are taking Topiramate GH

Things you must do

- Always follow your doctor's instructions carefully.
- Tell your doctor if you become pregnant while taking Topiramate GH.
- If you are about to start taking a new medicine, tell your doctor and pharmacist that you are taking Topiramate GH.
- Drink plenty of water. Topiramate GH has been known to cause kidney stones and drinking water may help prevent this.

Things you must not do

- Do not stop taking Topiramate GH without your doctor's permission.
- Do not take Topiramate GH to treat any other complaints unless your doctor says so.
- Do not give your medicine to anyone else, even if their symptoms seem similar to yours.
- Do not drink alcohol.
- Do not drive or operate machinery until you know how the medicine affects you.

Things to be careful of

Effects on thoughts and behaviour

Medicines used to treat epilepsy can increase the risk of suicidal thoughts and behaviour. If you experience feelings of deep sadness and unworthiness (depression) or a worsening of these feelings, any unusual changes in your mood or

the emergence of suicidal thoughts, behaviour or thoughts of self harm, you should report this to your doctor immediately.

Effects on driving and operating machinery

Topiramate GH may cause drowsiness, dizziness or other symptoms which could affect your ability to drive or operate machinery. It may also cause visual disturbances and/or blurred vision. Make sure you know how you are affected by this medicine before you drive or use machinery.

Particular care is recommended when you first start taking Topiramate GH or if the amount of Topiramate GH or any other medicine you are taking is increased or decreased.

Effects of food and alcohol

Topiramate GH can be taken with or without food.

Do not drink alcohol while taking Topiramate GH. Alcohol may increase the risk of unwanted side effects, such as drowsiness.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Topiramate GH.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Do not be alarmed by the following lists of side effects. You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- unusual tiredness, drowsiness, weakness, dizziness;
- decrease in appetite or weight loss;

- itchy skin or skin rash;
- inability to sleep;
- headache;
- tingling and numbness of hands and feet;
- nose bleed;
- nausea, diarrhoea, constipation;
- taste disturbance or loss of taste;
- decreased or lack of sweating.

Tell your doctor as soon as possible if you notice any of the following as you may need medical attention:

- nervousness or feeling anxious;
- disturbance in attention;
- difficulty with memory or memory impairment;
- slowing of thought processes;
- abnormal behaviour;
- balance disorder;
- co-ordination problems;
- expressive language disorder;
- unusual hair loss or thinning;
- abnormal frequent urination;
- decreased feeling or sensitivity, especially in the skin;
- fever or high temperature;
- unusual weakness;
- difficulty in speaking.

Tell your doctor immediately if you have:

- kidney stones;
- pain when passing urine;
- sudden changes in your eyesight (eg. blurred or loss of vision) or rapid uncontrollable movements of the eyes
- eye pain or increased pressure in eye.

These are serious side effects that may require urgent medical attention. Serious side effects are rare.

Tell your doctor or pharmacist if you notice anything that is making you feel unwell.

Other side effects not listed above may also occur in some people.

After taking Topiramate GH

Storage

Keep your medicine in the original container until you need to take it.

If you take it out of its original container it may not keep well.

Keep your medicine in a cool dry place where the temperature stays below 25°C.

Do not store Topiramate GH or any other medicine in the bathroom or near a sink. Do not leave it on a window sill or in the car.

Heat and dampness can destroy some medicines.

Keep it where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.

Product description

What it looks like

Topiramate GH tablets are available in bottles of 60 tablets:

- 25 mg tablets are white, round shaped, film-coated tablets, plain on both sides.
- 50 mg tablets are yellow, round shaped, film-coated tablets, plain on both sides.
- 100 mg tablets are white, round shaped, film-coated tablets, plain on both sides.
- 200 mg tablets are yellow, round shaped, film-coated tablets, plain on both sides.

Ingredients

Active Ingredient: topiramate.

Inactive Ingredients:

- lactose
- pregelatinised maize starch
- microcrystalline cellulose
- sodium starch glycollate
- magnesium stearate
- titanium dioxide
- hypromellose
- macrogol 400
- polysorbate 80
- iron oxide yellow (50mg and 200mg tablets only).

This medicine does not contain sucrose, gluten, tartrazine or any azo dyes.

Australian Registration Numbers

Topiramate GH 25mg tablets:
AUST R 135766

Topiramate GH 50mg tablets:
AUST R 135790

Topiramate GH 100mg tablets:
AUST R 135797

Topiramate GH 200mg tablets:
AUST R 135815

Distributor

Generic Health Pty Ltd
Level 1, 1102 Toorak Road
Camberwell VIC 3124

This leaflet was prepared in February 2015.