

GENVOYA Tablets

(150 mg elvitegravir / 150 mg cobicistat / 200 mg emtricitabine / 10 mg tenofovir alafenamide)

elvitegravir / cobicistat / emtricitabine / tenofovir alafenamide

Consumer Medicine Information

What is in this leaflet

Read all of this leaflet carefully before you start taking this medicine.

This leaflet answers some of the common questions about GENVOYA tablets. It does not contain all of the available information.

It does not take the place of talking to your doctor or pharmacist about your medical condition or treatment. If you have further questions, please ask your doctor or your pharmacist.

This medicine has been prescribed for you personally and you should not pass it on to others.

It may harm them, even if their symptoms are the same as yours.

Medicines are sometimes prescribed for conditions that are not mentioned in this leaflet.

Keep this leaflet with your GENVOYA medicine.

You may need to read it again.

What is GENVOYA

How GENVOYA works

GENVOYA tablets consist of the following four medicines:

- elvitegravir
- cobicistat
- emtricitabine
- tenofovir alafenamide

These are combined in one tablet to help control Human Immunodeficiency Virus (HIV) infection.

GENVOYA helps block HIV-1 reverse transcriptase, a viral chemical in your body (enzyme) that is needed for HIV-1 to multiply.

GENVOYA lowers the amount of HIV in the blood (viral load). GENVOYA may also help to increase the number of T cells (CD4⁺ cells), allowing your immune system to improve. Lowering the amount of HIV in the blood lowers the chance of death or infections that happen when your immune system is weak (opportunistic infections).

HIV infection destroys CD4 T cells, which are important to the immune system. The immune system helps fight infection. After a large number of T cells are destroyed, acquired immune deficiency syndrome (AIDS) may develop.

GENVOYA is for people who do not have a resistant HIV virus to GENVOYA.

Elvitegravir belongs to a class of antiviral medicines known as integrase inhibitors.

Cobicistat is a “booster”, to help increase the levels of elvitegravir.

Emtricitabine and tenofovir alafenamide belong to a group of antiviral medicines known as nucleoside and nucleotide reverse transcriptase inhibitors (NRTI).

Elvitegravir, cobicistat and emtricitabine are also components of STRIBILD® tablets.

Use in children

GENVOYA is used to treat HIV-1 infection in adults and children 12 years of age and older.

GENVOYA has not been studied in children under the age of 12 or weighing less than 35 kg, or adults over the age of 65.

Does GENVOYA cure HIV or AIDS

GENVOYA does not cure HIV infection or AIDS.

The long-term effects of GENVOYA are not known at this time.

People taking GENVOYA may still get opportunistic infections or other conditions that happen with HIV infection.

Opportunistic infections are infections that develop because the immune system is weakened. Some of these conditions are pneumonia, herpes virus infections, and *Mycobacterium avium complex* (MAC) infection.

This medicine is only available from a pharmacist after it has been prescribed by a doctor who specialises in the treatment of HIV infection.

If you wish to continue receiving treatment with GENVOYA it is

important you remain under the care of a hospital or doctor who specialises in the treatment of HIV infection.

Does GENVOYA reduce the risk of passing HIV to others

It is still possible to pass on HIV to other people through sexual contact, sharing needles, or being exposed to your blood when taking GENVOYA, although the risk is much lower with effective antiretroviral therapy.

Discuss with your doctor the precautions needed to avoid infecting other people. For your health and the health of others, it is important to always practice safer sex by using a latex or polyurethane condom or other barrier to lower the chance of sexual contact with semen, vaginal secretions, or blood.

Never re-use or share needles.

Before you take GENVOYA

When you must not take it

Together with your doctor, you need to decide whether GENVOYA is right for you. There are complex interactions with many different medications, so you should consult your doctor or pharmacist before starting any new medication.

Do not take GENVOYA if you are allergic to:

- elvitegravir,
- cobicistat,
- emtricitabine,
- tenofovir,
- or any of the other ingredients of GENVOYA.

The ingredients of GENVOYA are listed in the product description section of this leaflet.

Do not take GENVOYA if you are already taking any other

medicines that contain the same active ingredients such as:

- ATRIPLA®
- EMTRIVA®
- TRUVADA®
- EVIPLERA®
- STRIBILD®
- TYBOST®
- ODEFSEY®
- DESCOVY®

Do not take GENVOYA if you are taking other medicines that contain:

- lamivudine (e.g. Combivir, Zeffix, Kivexa, Trizivir, Triumeq)
- ritonavir (e.g. Kaletra)
- tenofovir disoproxil fumarate (e.g. VIREAD)
- Efavirenz (e.g. Stocrin)

These medicines contain similar active ingredients as GENVOYA.

Do not take GENVOYA if you take:

- alfuzosin hydrochloride (e.g. Xatral),
- carbamazepine (e.g. Tegretol),
- phenobarbital or phenytoin (e.g. Dilantin),
- ergot-containing medicines like dihydroergotamine, ergotamine (e.g. Cafergot, Dihydergot, Migerot).
- lovastatin (e.g. Mevacor)
- midazolam (e.g. Hypnovel)
- rifabutin (e.g. Mycobutin)
- sildenafil (e.g. Viagra/Revatio)
- simvastatin (e.g. Invast/Zimcol)
- tadalafil (e.g. Cialis/Adcirca)
- triazolam (e.g. Halcion)
- rifampicin (e.g. Rifadin/Rimycin)
- St John's Wort or products containing St John's Wort.

Do not take GENVOYA to treat your HIV infection if you are also taking adefovir dipivoxil to treat your hepatitis B virus (HBV) infection.

This is not a complete list of medicines that you should not take with GENVOYA. You

should discuss this with your doctor.

Before you start to take it

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if you have, or have had, any of the following medical conditions:

- severe kidney problems or are undergoing kidney dialysis treatment.
- liver problems, including hepatitis B or C virus infection.

Tell your doctor if you take an antacid medicine that contains aluminium, magnesium hydroxide, or calcium carbonate.

Take antacids at least 2 hours before or after you take GENVOYA.

Tell your doctor if you are pregnant, or likely to become pregnant during your course of medication.

We do not know if GENVOYA can harm your unborn child. You and your doctor will need to decide if GENVOYA is right for you.

Tell your doctor if you are breastfeeding, or likely to breastfeed during your course of medication.

You should not breastfeed if you are HIV-positive because of the chance of passing the HIV virus to your baby. One of the active substances in this medicine (emtricitabine) has been found in breast milk at low concentrations. Talk with your doctor about the best way to feed your baby.

Taking other medicines

Some medicines may affect the levels of GENVOYA or GENVOYA may affect the levels of other medicines in the body when they are taken at the same time as GENVOYA.

Your doctor may change your other medicines or change their doses. Other medicines, including herbal products may affect GENVOYA.

For this reason, it is very important to let your doctor or pharmacist know what medications, herbal supplements, or vitamins you are taking.

Know the medicines you take. Keep a list of medicines and show it to your doctor and pharmacist when you get a new medicine.

Your doctor and your pharmacist can tell you if you can take these medicines with GENVOYA.

Do not start any new medicines while you are taking GENVOYA without first talking with your doctor or pharmacist.

How to take GENVOYA

Take the exact amount of GENVOYA your doctor has prescribed for you.

Never change the dose on your own.

Do not stop this medicine unless your healthcare provider tells you to stop.

How much to take

The usual dose is one GENVOYA tablet orally, once daily.

Always take GENVOYA with food.

If you forget to take it

Do not miss a dose of GENVOYA.

If you forget to take GENVOYA, take your missed dose right away unless it is almost time for your next dose.

Do not take a double dose to make up for a forgotten dose.

Continue with your regular dosing schedule.

When your GENVOYA supply starts to run low, get more from your doctor or pharmacy.

This is very important because the amount of virus in your blood may increase if the medicine is stopped for even a short time. The virus may develop resistance to GENVOYA and become harder to treat.

If you take too much (overdose)

Immediately telephone your doctor or Poisons Information Centre: 131126 (Australia) and 0800 764 766 (New Zealand) or go to the Accident and Emergency department at your nearest hospital if you think you or anyone else may have taken too many GENVOYA tablets. Do this even if there are no signs of discomfort or poisoning. This may need urgent medical attention.

While you are taking GENVOYA

Things you must not do

Do not breastfeed. See “Before you start to take it”

Avoid doing things that can spread HIV infection since GENVOYA does not stop you from passing the HIV Infection to others.

- Do not share needles or other injection equipment.
- Do not share personal items that can have blood or body fluids on them, like toothbrushes or razor blades.
- Do not have any kind of sex without protection.

Always practice safer sex by using a latex or polyurethane condom or other barrier to reduce the chance of sexual contact with semen, vaginal secretions, or blood.

Do not take GENVOYA after the expiry or “use by” date (EXP) printed on the bottle.

If you take it after the expiry date has passed, it may not work as well.

Do not take GENVOYA if the packaging is torn or shows signs of tampering.

Things to be careful of

Be careful driving or operating machinery until you know how GENVOYA affects you.

If you are dizzy, have trouble concentrating, or are drowsy, avoid activities that may be dangerous, such as driving or operating machinery.

Side Effects

Like all medicines, GENVOYA can have side effects, although not everybody gets them. Some may be serious and need medical attention.

Check with your doctor as soon as possible if you have any problems while taking GENVOYA, even if you do not think the problems are connected with the medicine or are not listed in this leaflet.

Serious side effects

Lactic Acidosis

If you have any of the following symptoms after taking your medication, tell your doctor IMMEDIATELY or go to the accident and emergency department at your nearest hospital.

- you feel very weak or tired
- you have unusual (not normal) muscle pain
- you have trouble breathing
- you have stomach pain with nausea and vomiting
- you feel cold, especially in your arms and legs
- you feel dizzy or light headed
- you have a fast or irregular heartbeat

These side effects may be due to a condition called lactic acidosis (build-up of an acid in the blood).

Lactic acidosis can be a medical emergency and may need to be treated in the hospital.

Serious Liver Problems (hepatotoxicity)

If you have any of the following symptoms while taking your medication, tell your doctor IMMEDIATELY or go to the accident and emergency department at your nearest hospital.

- your skin or the white part of your eyes turns yellow (jaundice)
- your urine turns dark
- your bowel movements (stools) turn light in colour
- you don't feel like eating food for several days or longer
- you feel sick to your stomach (nausea)
- you have lower stomach area (abdominal) pain

These side effects may be due to a condition called hepatotoxicity with liver enlargement (hepatomegaly) and fat deposits in the liver (steatosis) which sometimes occurs in patients taking anti-HIV medicines.

You may be more likely to get lactic acidosis or liver problems if you are female, very overweight (obese), or have been taking similar nucleoside analog-containing medicines, like GENVOYA, for a long time.

Hepatic Flares

If you have HIV infection and HBV infection you should not stop your GENVOYA treatment without first discussing this with your doctor.

Your HBV may get worse (flare-up) if you stop taking GENVOYA. A "flare-up" or "hepatic flare" is when your HBV infection suddenly returns in a worse way than before. You may require medical exams and blood tests for several months after stopping treatment.

GENVOYA is not approved for the treatment of HBV, so you must discuss your HBV therapy with your doctor.

Changes in body fat

Changes in body fat develop in some people receiving anti HIV-1 therapy. These changes may include an increased amount of fat in the upper back and neck ('buffalo hump'), in the breasts and around the trunk. Loss of fat from the legs, arms and face may also happen. The cause and long-term health effects of these fat changes are not known.

Signs and symptoms of inflammation

In some patients with advanced HIV infection (AIDS), signs and symptoms of inflammation from previous infections may occur soon after anti-HIV treatment is started. It is believed that these symptoms are due to an improvement in the body's immune response, which lets the body fight infections that may have been present with no obvious symptoms. If you notice any symptoms of infection, please tell your doctor immediately.

Allergy

Some people are allergic to medicines. If you have any of the following symptoms soon after taking your medicine, DO NOT TAKE ANY MORE GENVOYA and tell your doctor IMMEDIATELY or go to the accident and emergency department at your nearest hospital:

- skin troubles such as lumpy skin rash or "hives"
- swelling of the face, lips, mouth or throat which may cause difficulty in swallowing or breathing
- wheezing, chest pain or tightness
- fainting

These are very serious effects. If you have them, you may have a serious allergic reaction. You may need urgent medical attention or hospitalisation. Hypersensitivity reactions are very rare.

Common side effects

The most common side effect of GENVOYA is nausea.

Other side effects include:

- diarrhea
- fatigue
- headaches
- abdominal pain
- indigestion
- flatulence
- rash
- vomiting

Talk to your doctor or pharmacist if you don't understand anything in this list.

This is not a complete list of side effects possible with GENVOYA.

Ask your doctor or pharmacist for a more complete list of side effects of GENVOYA and all the medicines you will take.

After taking GENVOYA

Storage

Keep GENVOYA tablets where children cannot reach them.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store them.

Keep GENVOYA tablets in a cool, dry place where it stays below 25°C.

Do not store GENVOYA or any other medicine in a bathroom or near a sink.

Do not leave GENVOYA in the car or on a window sill.

Heat and dampness can destroy some medicines.

Keep your GENVOYA tablets in the bottle with the cap tightly closed until you take them.

If you take GENVOYA tablets out of their pack they may not keep well.

Product Description

What the tablets look like

GENVOYA tablets are capsule-shaped, film-coated and green in colour.

Each tablet is debossed with "GSI" on one side and the number "510" on the other side.

GENVOYA tablets are supplied in bottles containing 30 tablets.

Ingredients

Each GENVOYA tablet contains the active ingredients:

- elvitegravir
- cobicistat
- emtricitabine
- tenofovir alafenamide

Each GENVOYA tablet also contains the following inactive ingredients:

- lactose monohydrate
- microcrystalline cellulose
- croscarmellose sodium
- hypromellose
- silicon dioxide
- sodium lauryl sulfate
- magnesium stearate

Film-coating:

- polyvinyl alcohol
- titanium dioxide
- polyethylene glycol
- talc
- indigo carmine aluminum lake
- iron oxide yellow

Sponsor

GENVOYA tablets are supplied in Australia by:

Gilead Sciences Pty Ltd
Level 6, 417 St Kilda Road
Melbourne, Victoria 3004

In New Zealand:

Gilead Sciences (NZ)
c/- PricewaterhouseCoopers
Level 8 Pricewaterhousecoopers
Tower
188 Quay Street
Auckland 1010

Date of preparation: 10 November 2016

AUST R 233398

GENVOYA, STRIBILD, EMTRIVA, TRUVADA, EVIPLERA, ODEFSEY, DESCOVY, TYBOST, VIREAD and GSI are trademarks of Gilead Sciences, Inc. or one of its related companies. ATRIPLA is a registered trademark of Bristol-Myers Squibb & Gilead Sciences, LLC. All other trademarks referenced herein are the property of their respective owners.