

TELMISARTAN HCT GH Tablets

Telmisartan and Hydrochlorothiazide

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about TELMISARTAN HCT GH.

It does not contain all available information. It does not take the place of talking to your doctor or pharmacist.

This leaflet was last updated on the date at the end of this leaflet. More recent information may be available.

The latest Consumer Medicine Information is available from your pharmacist, doctor, or from www.medicines.org.au and may contain important information about the medicine and its use of which you should be aware.

All medicines have benefits and risks. Your doctor has weighed the risks of you taking this medicine against the benefits it is expected to have for you.

If you have any concerns about this medicine, ask your doctor or pharmacist. Keep this leaflet with the medicine.

You may need to read it again.

What TELMISARTAN HCT GH is used for

TELMISARTAN HCT GH is used to lower high blood pressure (hypertension).

Everyone has blood pressure. This pressure helps your blood move around your body. Your blood pressure may be different at different times of the day,

depending on how busy or worried you are.

You have hypertension (high blood pressure) when your blood pressure stays higher than is needed, even when you are calm or relaxed. There are usually no signs of hypertension. The only way of knowing that you have hypertension is to have your blood pressure checked on a regular basis.

If high blood pressure is not treated, it can lead to serious health problems, including stroke, heart disease and kidney failure.

How TELMISARTAN HCT GH works

TELMISARTAN HCT GH contains telmisartan and hydrochlorothiazide in one single tablet. These two active ingredients help to reduce blood pressure in different ways:

- Telmisartan belongs to a group of medicines called angiotensin II receptor antagonists. Angiotensin II is a substance in the body which causes blood vessels to narrow, thus increasing blood pressure. Telmisartan works by blocking the effect of angiotensin II. When the effect of angiotensin II is blocked, your blood vessels relax and your blood pressure goes down.
- Hydrochlorothiazide belongs to the group of medicines called diuretics. Diuretics help to reduce the amount of excess fluid in the body by increasing the amount of urine produced. They help with lowering blood pressure particularly when used with other blood pressure

lowering medicines. Your doctor may have prescribed TELMISARTAN HCT GH for another reason.

Ask your doctor if you have any questions about why TELMISARTAN HCT GH has been prescribed for you.

TELMISARTAN HCT GH is not addictive. This medicine is available only with a doctor's prescription.

Use in Children

The safety and effectiveness of TELMISARTAN HCT GH in children and teenagers up to 18 years of age have not been established.

Before you take TELMISARTAN HCT GH

When you must not take it

Do not take TELMISARTAN HCT GH if you have ever had an allergic reaction after taking:

- Telmisartan or hydrochlorothiazide (the two active ingredients in TELMISARTAN HCT GH),
- other sulfonamide-derived medicines
- any of the other ingredients listed at the end of this leaflet.

Symptoms of an allergic reaction to TELMISARTAN HCT GH may include skin rash, itchiness, shortness of breath, swelling of the face, lips or tongue, muscle pain or tenderness or joint pain.

Do not take TELMISARTAN HCT GH if you have a rare hereditary condition of fructose intolerance.

The maximum recommended daily dose of TELMISARTAN HCT GH contains 169 mg of sorbitol (in the 40/12.5 mg tablets) and approximately 338 mg of sorbitol (in the 80/12.5 mg and 80/25 mg tablets).

Do not take TELMISARTAN HCT GH if you have a rare hereditary condition of galactose intolerance.

The maximum recommended daily dose of TELMISARTAN HCT GH contains 112 mg of lactose (in the 40/12.5 mg and 80/12.5 mg tablets) and 99 mg of lactose (in the 80/25 mg tablets).

Do not take TELMISARTAN HCT GH if you are pregnant or intend to become pregnant.

TELMISARTAN HCT GH is not recommended for use in pregnancy. Like other similar medicines, it could affect your unborn baby if you take it during pregnancy.

Do not breast-feed if you are taking TELMISARTAN HCT GH.

It is not known if telmisartan or hydrochlorothiazide, the active ingredients in TELMISARTAN HCT GH, pass into breast milk and there is a possibility that your baby may be affected.

Do not take TELMISARTAN HCT GH if you have any of the following medical conditions:

- severe liver disease
- severe kidney disease
- cholestasis or biliary obstructive disorders (problem with the flow of bile from the gall bladder)
- low potassium levels in the blood
- high calcium levels in the blood.

Do not take TELMISARTAN HCT GH after the expiry date printed on the pack.

Do not take TELMISARTAN HCT GH if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should start taking TELMISARTAN HCT GH, talk to your doctor.

Before you start to take it

Tell your doctor if you have allergies to:

- any other medicines
- any other substances, such as foods, preservatives or dyes

Tell your doctor if you have, or have had, any medical conditions, especially the following:

- kidney problems
- liver problems
- heart problems
- a condition known as primary hyperaldosteronism (raised aldosterone levels, also known as Conn's syndrome)
- diabetes
- gout
- fructose intolerance
- galactose intolerance
- a low salt diet
- recent severe diarrhoea or vomiting
- asthma
- systemic lupus erythematosus (a disease affecting the skin, joints and kidney)

Tell your doctor if you are following a very low salt diet.

Tell your doctor if you plan to become pregnant or breastfeed.

If you have not told your doctor about any of the above, tell your doctor before you take TELMISARTAN HCT GH .

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and TELMISARTAN HCT GH may interfere with each other. These include:

- any other medicines used to treat high blood pressure such as beta blockers
- heart medicines such as digoxin, a medicine used to treat heart failure or antiarrhythmic medicines
- lithium, a medicine used to treat certain mental illnesses
- other diuretics or fluid tablets medicines used to help the kidneys get rid of salt and water by increasing the amount of urine produced
- laxatives
- potassium supplements or potassium-containing salt substitutes
- medicines or salt substitutes which may increase your potassium levels
- alcohol
- sleeping tablets
- strong pain killing medicines
- medicines for diabetes (oral tablets or capsules or insulin)
- powder or granules used to help reduce cholesterol
- corticosteroid medicines such as prednisolone, cortisone or ACTH
- aspirin
- nonsteroidal anti-inflammatory agents (medicines used to relieve pain, swelling and other symptoms of inflammation including arthritis)
- medicines used to treat gout
- calcium supplements or medicines containing calcium
- anticholinergic medicines, which can be used to treat Parkinson's disease, relieve stomach cramps or prevent travel sickness
- amantadine, a medicine used to treat Parkinson's disease or to prevent influenza
- medicines used to treat cancer (cytotoxic medicines)

These medicines may be affected by TELMISARTAN HCT GH, or may affect the way it works. Also, other medicines used to treat high blood pressure may have an additive effect with TELMISARTAN HCT GH in lowering your blood pressure. Therefore, you may need different amounts of your medicines.

Your doctor or pharmacist may have more information on medicines to be careful with or avoid while taking TELMISARTAN HCT GH.

How to take TELMISARTAN HCT GH

Follow all directions given to you by your doctor or pharmacist carefully.

They may differ from the information contained in this leaflet. Your doctor or pharmacist will tell you how many tablets you will need to take each day. This depends on your condition and whether or not you are taking any other medicines.

If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

How much to take

The usual dose for adults is one TELMISARTAN HCT GH 40/12.5 mg tablet once a day. If your blood pressure is still too high after 4–8 weeks of starting treatment, your doctor may increase your dose to one TELMISARTAN HCT GH 80/12.5 mg tablet once a day.

If your blood pressure is still not satisfactorily controlled with TELMISARTAN HCT GH 80/12.5 mg, your doctor may increase your dose to one TELMISARTAN HCT GH 80/25 mg tablet once a day.

It is important to take TELMISARTAN HCT GH exactly as your doctor or pharmacist has told you.

When to take it

Take TELMISARTAN HCT GH at about the same time each day, either morning or evening.

Taking it at the same time each day will have the best effect. It will also help you remember when to take it.

It does not matter if you take TELMISARTAN HCT GH before or after food.

How to take it

Swallow the tablet whole with a full glass of water.

You can take TELMISARTAN HCT GH with or without food.

How long to take it

Take TELMISARTAN HCT GH every day until your doctor tells you to stop.

TELMISARTAN HCT GH helps control your high blood pressure but does not cure it. It is important to keep taking TELMISARTAN HCT GH every day even if you feel well.

People who have high blood pressure often feel well and do not notice any signs of this problem.

If you forget to take it

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to. Otherwise, take the dose as soon as you remember, and then go back to taking it as you would normally.

Do not take a double dose to make up for the dose that you missed.

This may increase the chance of you getting unwanted side effects.

If you have trouble remembering when to take your medicine, ask your pharmacist for some hints.

If you take too much (overdose)

Immediately telephone your doctor or Poisons Information Centre (telephone 13 11 26) for advice or go to Accident and Emergency at your nearest

hospital, if you think that you or anyone else may have taken too much TELMISARTAN HCT GH.

Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention. If you take too much TELMISARTAN HCT GH you may feel dizzy, light-headed or faint. Your heartbeat may be faster or lower than usual. You may experience dehydration, nausea, drowsiness and muscle spasm.

While you are taking TELMISARTAN HCT GH

Things you must do

Tell any other doctors, dentists and pharmacists who are treating you that you are taking TELMISARTAN HCT GH.

If you are about to be started on any new medicine, tell your doctor or pharmacist that you are taking TELMISARTAN HCT GH.

If you feel that TELMISARTAN HCT GH is not helping your condition, tell your doctor or pharmacist.

Tell your doctor if, for any reason, you have not used TELMISARTAN HCT GH exactly as prescribed.

Otherwise, your doctor may think that it was not effective and change your treatment unnecessarily.

Tell your doctor immediately if you become pregnant while taking TELMISARTAN HCT GH.

You should not take this medicine while you are pregnant.

If you are going to have surgery, tell your doctor and anaesthetist that you are taking TELMISARTAN HCT GH.

TELMISARTAN HCT GH may affect some medicines you receive during surgery.

Things you must not do

Do not use TELMISARTAN HCT GH to treat other complaints unless your doctor or pharmacist tells you to.

Do not give this medicine to anyone else, even if they have the same symptoms as you.

Things to be careful of

Be careful when driving or operating machinery until you know how TELMISARTAN HCT GH affects you.

Like other medicines used to treat high blood pressure, TELMISARTAN HCT GH may cause sleepiness, dizziness or lightheadedness in some people.

If you have any of these symptoms, do not drive, operate machinery or do anything else that could be dangerous.

You may feel dizzy or light-headed when you begin to take TELMISARTAN HCT GH, especially if you are also taking a diuretic (or fluid tablet) or if you are dehydrated.

If this medicine makes you feel dizzy or light-headed, be careful when getting up from a sitting or lying position.

Standing up slowly, especially when you get up from a bed or chair, will help your body get used to the change in position and blood pressure. If this problem continues or gets worse, talk to your doctor.

If you exercise, or if you sweat, or if the weather is hot, you should drink plenty of water.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking TELMISARTAN HCT GH even if you do not think it is connected with the medicine.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need

medical treatment if you get some of the side effects.

Do not be alarmed by this list of possible side effects.

You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have. Tell your doctor if you notice any of the following and they worry you:

- ‘flu-like’ symptoms
- fainting, dizziness or spinning sensation
- a feeling of tension or fullness in the nose, cheeks and behind the eyes, sometimes with a throbbing ache (sinusitis)
- infections of the air passages
- shortness of breath or difficulty breathing
- abnormal or blurred vision
- back pain
- changes in heart rhythm or increased heart rate
- rash or redness or itchiness of skin
- increased sweating
- dizziness or lightheadedness when you stand up (postural hypotension)
- stomach pain or discomfort (abdominal pain, dyspepsia, gastritis)
- wind or excessive gas in the stomach or bowel
- vomiting
- diarrhoea or constipation
- dry mouth
- pins and needles
- sleep disturbances or trouble sleeping
- feeling anxious
- depression
- impotence
- leg pain or cramps in legs
- aching muscles or aching joints not caused by exercise or muscle spasms
- chest pain
- pain
- liver problems

- changes in the levels of potassium or sodium or uric acid in your blood (such changes are usually detected by a blood test)

Tell your doctor as soon as possible if you experience any side effects during or after taking TELMISARTAN HCT GH, so that these may be properly treated.

Symptoms such as feeling very thirsty, sleepy, sick or vomiting, a dry mouth, general weakness, muscle pain or cramps, a very fast heart rate, may mean that the hydrochlorothiazide part of TELMISARTAN HCT GH is having an excessive effect.

You should tell your doctor if you experience any of these symptoms. Tell your doctor immediately or go to casualty at your nearest hospital if you notice any of the following:

- swelling of the face, lips, mouth, tongue or throat which may cause difficulty in swallowing or breathing
- severe and sudden onset of pinkish, itchy swellings on the skin, also called hives or nettle rash
- developing or worsening of a disease called systemic lupus erythematosus which affects the skin, joints and kidney

These are serious side effects. You may need urgent medical attention or hospitalisation. These side effects are rare.

Tell your doctor if you notice anything else that is making you unwell.

Other side effects not listed above may also occur in some people.

After taking TELMISARTAN HCT GH

Storage

Leave the tablets in the blister strip until it is time to take a dose.

The blister pack protects the tablets from light and moisture.

Keep TELMISARTAN HCT GH in a cool, dry place where the temperature stays below 25°C.

Do not store TELMISARTAN HCT GH or any other medicine in the bathroom or near a sink. Do not leave it in the car or on window sills.

Heat and dampness can destroy some medicines.

Keep TELMISARTAN HCT GH where young children cannot reach it.

Disposal

If your doctor tells you to stop using TELMISARTAN HCT GH or it has passed its expiry date, ask your pharmacist what to do with any that is left over.

Product description

What it looks like

TELMISARTAN HCT GH is the brand name of your medicine. TELMISARTAN HCT GH tablets are available in three strengths: 40/12.5 mg, 80/12.5 mg and 80/25 mg tablets.

TELMISARTAN HCT GH 40/12.5 mg and 80/12.5 mg tablets are pink mottled and white to off-white biconvex, oval shaped, two layer tablets. The white to off-white layer may contain pink specks. Each tablet contains either 40 mg or 80 mg of telmisartan and 12.5 mg of hydrochlorothiazide. The pink mottled face of TELMISARTAN HCT GH 40/12.5 mg tablets are marked with L199. The pink mottled face of TELMISARTAN

HCT GH 80/12.5 mg tablets are marked with L200.

TELMISARTAN HCT GH 80/25 mg tablets are yellow mottled and white to off-white biconvex, oval shaped, two layer tablets. The white to off-white layer may contain yellow specks. Each tablet contains 80 mg of telmisartan and 25 mg of hydrochlorothiazide. The white face of TELMISARTAN HCT GH 80/25 mg tablets are marked with L201.

TELMISARTAN HCT GH tablets are available in blister packs of 28 tablets.

The following Australian Registration Numbers appear on the carton:

AUST R 208070 for
TELMISARTAN HCT GH 40/12.5 mg tablets

AUST R 208075 for
TELMISARTAN HCT GH 80/12.5 mg tablets

AUST R 208076 for
TELMISARTAN HCT GH 80/25 mg tablets

Ingredients

Each TELMISARTAN HCT GH 40/12.5 mg tablet contains 40 mg telmisartan and 12.5 mg hydrochlorothiazide.

Each TELMISARTAN HCT GH 80/12.5 mg tablet contains 80 mg telmisartan and 12.5 mg hydrochlorothiazide.

Each TELMISARTAN HCT GH 80/25 mg tablet contains 80 mg telmisartan and 25 mg hydrochlorothiazide.

The other ingredients found in the tablets are:

- povidone (K25)
- lactose
- magnesium stearate
- meglumine
- sodium hydroxide
- sodium stearyl fumarate
- mannitol

TELMISARTAN HCT GH 40/12.5 mg and 80/12.5 mg tablets also contain Pigment Blend PB-24880 Pink and TELMISARTAN HCT GH 80/25 mg tablets also contain Pigment Blend PB-52290 Yellow, as colouring agent.

Sponsor

GENERIC HEALTH PTY LTD
Level 1, 1102 Toorak Road
Camberwell VIC 3124

This leaflet was last updated in September 2015