

Canesten® Clotrimazole Thrush Treatment Once Pessary

Clotrimazole

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Canesten Thrush Treatment.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your pharmacist has weighed the risks of you using Canesten Thrush Treatment against the benefits they expect it will have for you.

If you have any concerns about using this treatment, ask your doctor or pharmacist.

Keep this leaflet with the medicine.
You may need to read it again.

What Canesten Thrush Treatment is used for

Canesten Thrush Treatment is used to treat a fungal infection in the vagina and vulval area (thrush).

This medicine belongs to a group of medicines called azole antifungals.

What is vaginal thrush?

Candida albicans is a yeast fungus that can multiply rapidly and cause damage to the vagina when other micro-organisms aren't present in a normal balance.

Some situations which can increase susceptibility to thrush are:

- Diabetes
- Obesity
- Illness
- Fatigue
- Hormonal fluctuations (pregnancy, menstrual cycle, oral contraceptive pill, HRT)
- Prolonged use of antibiotics or steroids

Thrush may cause the following symptoms:

- Vaginal itching
- Vaginal soreness
- A white odourless discharge from the vagina (like cottage cheese)
- Pain during intercourse

The symptoms below are not usually signs of thrush:

- Abnormal or irregular vaginal bleeding
- Blood stained discharge
- Vulva or vaginal sores, ulcers or blisters
- Lower abdominal pain
- Pain when passing urine
- Fever or chills
- Foul smelling and/or unusual colour discharge

See your doctor if you experience any of the above symptoms.

Ask your doctor or pharmacist if you have any questions about whether this medicine is suitable for you.

This medicine is available without a doctor's prescription, only through your pharmacist.

Before you use Canesten Thrush Treatment

When you must not use it

Do not use Canesten Thrush Treatment if you have an allergy to:

- any medicine containing clotrimazole
- any of the ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include swelling to the area to which treatment has been applied; rash, itching or hives on the skin.

Do not use this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should start using this medicine, talk to your doctor or pharmacist.

Before you start to use it

See your doctor if:

- this is your first infection
- you are under 18 years of age
- you have had 3 or more infections in the last 6 months
- you have allergies to any other topical treatments.

Tell your doctor if you are pregnant.

You should see your doctor if you suspect you have thrush in the first three months of your pregnancy. Your doctor can discuss with you the risks and benefits involved with treatment.

Tell your doctor if you are breast-feeding and want to use this treatment.

It is not known whether the active ingredient, clotrimazole, is excreted in breast milk. Your doctor will be able to discuss with you the risks and benefits involved with treatment.

Tell your doctor if you are diabetic.

Thrush in diabetics can mean that your blood sugar levels are not well controlled. Your doctor will be able to check this for you and advise treatment.

If you have not told your doctor about any of the above, tell him/her before you start taking Canesten Thrush Treatment.

Taking other medicines

There are no reports of Canesten Thrush Treatment interfering with other medicines.

How to use Canesten Thrush Treatment

Follow all directions given to you by your doctor or pharmacist carefully.

They may differ from the information contained in this leaflet.

If you do not understand the instructions on the box, ask your doctor or pharmacist for help.

How to use it

Wash your hands thoroughly before handling the pessary or using the applicator.

To avoid the possibility of re-infection, strict personal hygiene must be observed.



The pessary should preferably be inserted in the evening, just before bed time, with the use of the applicator. Using it before you go to sleep will have the best effect.

Loading the applicator

1. Wash your hands thoroughly.
2. Pull out the plunger until it stops (A).
3. Remove the pessary from the blister. Squeeze the cup end of the applicator (B) between thumb and index finger to slightly flatten the cup.
4. Place the pessary into the flattened cup end of the applicator.

Inserting the applicator

5. Carefully insert the end of the applicator containing the pessary as gently and deeply into the vagina as possible. This is best achieved when lying on your back with your knees bent.
6. Hold the applicator in position and slowly push the plunger with the forefinger until it stops. This places the pessary in the vagina.
7. Remove the applicator from the vagina.
8. Discard the applicator.



How long to use it

Insert one pessary, once.

If you are not sure what to do, ask your doctor or pharmacist.

DO NOT TAKE THE PESSARY BY MOUTH

Seek medical advice if you or anyone else has taken Canesten Once pessary by mouth. Do this even if there are no signs of discomfort or poisoning.

You may need medical attention.

While you are using Canesten Thrush Treatment

Things you must do

Discontinue treatment and tell your doctor or pharmacist if you develop a local reaction to this product.

See your doctor if the symptoms are not gone within 4 days.

Things you must not do

Do not take Canesten Thrush Treatment to treat any other complaints unless your doctor tells you to.

Do not give your treatment to anyone else, even if you suspect they have the same condition as you.

Things to be careful of

Canesten Thrush Treatment may reduce the effectiveness and safety of latex products, such as condoms and diaphragms.

This effect is temporary and occurs only during treatment.

Things that may be helpful for resolving Thrush

Some self help measures suggested below may help your condition. Your doctor or pharmacist can give you more information about these measures.

- Blot with toilet paper after urinating (instead of wiping back to front).
- Change underwear, night attire, towels and linen daily.

Avoid where possible:

- perfumed toilet paper
- fragrant soaps and bath oils
- wearing tight or synthetic underwear
- douches or vaginal deodorants

Side effects

Tell your doctor or pharmacist if you experience discomfort while you are using Canesten Thrush Treatment.

This medicine helps most people with thrush but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Do not be alarmed by the following side effect.

You may not experience it.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- skin reaction to the pessary (burning, stinging, redness)

Any of the above may be signs of an allergic reaction. Your doctor or pharmacist may recommend you discontinue treatment if you experience these effects.

After using Canesten Thrush Treatment

Storage

Keep the pessary in the protective blister pack until it is time to use it.

Keep your pessary in a cool dry place where the temperature stays below 25°C.

Keep the treatment where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Expiry

The expiry date is marked on the end panel of the carton and is also printed on the back of the blister. The date refers to the month and year of expiration eg 0805, refers to the eighth month of 2005. The product can be used until the end of this month.

Disposal

If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.

Discard the medicine out of the reach of children.

Product description

What it looks like

Canesten Thrush Treatment Once Pessary is a white pellet-shaped pessary packaged in an aluminium blister.

The package carton contains a plastic applicator for the insertion of the pessary. The plastic applicator is for use with the Canesten pessary only.

Discard applicator after use.

Ingredients

Canesten Thrush Treatment Once Pessary contains 500 mg clotrimazole as the active ingredient. It also contains:

- lactose
- cellulose-microcrystalline
- lactic acid
- starch-maize
- crospovidone
- calcium lactate
- magnesium stearate
- silicon dioxide
- hypromellose

This medicine does not contain sucrose, gluten, tartrazine or any other azo dyes.

Manufacturer

Canesten® Thrush Treatment Once Pessary is made in Germany.

Canesten® Thrush Treatment is distributed in Australia by:

Bayer Australia Limited
(ABN 22 000 138 714)
Consumer Care Business Group
875 Pacific Highway
PYMBLE NSW 2073

Toll Free: 1800 023 884

Canesten® Thrush Treatment is distributed in New Zealand by:

Bayer New Zealand Limited
Consumer Care Business Group
Argus Place
Glenfield
AUCKLAND NZ

Toll Free: 0800 847 874

® = Registered Trademark

This leaflet was prepared in August 2005.