

# IMIGRAN® Mk II Injection

(Sumatriptan succinate)

## Consumer Medicine Information

### What is in this leaflet?

Please read this leaflet carefully before you start using *Imigran* Mk II injection.

This leaflet answers some common questions about *Imigran* Mk II injection. It does not contain all of the available information.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the expected benefits of you using *Imigran* Mk II injection against the risks this medicine could have for you.

**If you have any concerns about taking this medicine, ask your doctor or pharmacist.**

**Keep this leaflet with the medicine.**

You may need to read it again.

### What is *Imigran* Mk II injection used for?

*Imigran* Mk II injection contains the active ingredient sumatriptan succinate. This medicine belongs to a group of drugs called serotonin agonists.

*Imigran* Mk II injection is used to relieve a migraine attack or cluster headache. It should not be used to prevent migraine attacks or cluster headaches from occurring. *Imigran* Mk II injection may be used for migraine headaches with or without what is known as 'aura'.

It is thought that migraine headaches and cluster headaches are due to widening of certain blood vessels in the head. *Imigran* Mk II injection works by making those vessels normal again and eases the symptoms of migraine and cluster headaches.

Your *Imigran* Mk II injection does not work in other types of headache which are not a migraine or a cluster headache.

*Imigran* Mk II injection is not addictive.

### Before you use *Imigran* Mk II injection

*Do not use if:*

You must not use *Imigran* Mk II injection if:

- **you have ever had an allergic reaction to sumatriptan succinate (See "Side-Effects") or any of the ingredients listed toward the end of this leaflet. (See "Ingredients").**
- **you have or have had:**
  - heart disease or heart attack.
  - shortness of breath, pain or tightness in the chest, jaw or upper arm.
  - peripheral vascular disease (pain in the back of the legs) or are prone to cold, tingling or numb hands and feet.
  - Prinzmetal's angina (an uncommon form of angina where pain is experienced at rest rather than during activity).

- angina.
- high blood pressure.
- stroke.
- severe liver disease.

- **you have taken any of these medicines in the last 24 hours:**
  - Ergotamine (eg. Cafergot)
  - Dihydroergotamine (eg. Dihydergot)
  - Methysergide (eg. Deseril)
  - naratriptan (eg Naramig)
  - zolmitriptan (eg Zomig).
- **you have taken any of these medicines in the last two weeks:**
  - Monoamine oxidase inhibitors (MAOIs), a type of medicine used for depression.
  - SSRIs (Selective Serotonin Reuptake Inhibitors) or SNRIs (Serotonin-Noradrenaline Reuptake Inhibitors) used to treat depression.
- **the expiry date (EXP) printed on the pack has passed.**
- **the packaging is torn or shows signs of tampering.**

*Tell your doctor if:*

You must tell your doctor if:

- **you are allergic to foods, dyes, preservatives or any other medicines, including any that contain sulphur (eg. sulphonamide antibiotics).**
- **you are taking or have taken any other medicines in the last two weeks, including medicines you buy without a prescription, particularly herbal preparations containing St John's Wort and**

medicines prescribed for depression.

- you are breastfeeding, pregnant or trying to become pregnant.
- you have or have had medical conditions like:
  - liver or kidney problems.
  - heart problems. Risk factors including high blood pressure, even if it is under control, high blood cholesterol levels, a family history of heart problems, obesity, diabetes, you are male and over 40 years of age, you are female and have undergone menopause or you smoke.
  - epilepsy, seizures, or fits or been told that you are prone to this problem
  - stroke

---

## How do I use *Imigran* Mk II injection?

---

The first time you ever have an *Imigran* Mk II injection it must be under the supervision of a doctor. After this you must use the injection as your doctor has instructed.

See the patient instruction leaflet inside the pack for information on how to load the Autoinjector with the syringe and how to discard the empty syringes. If there is something you do not understand, ask your doctor or pharmacist.

### *How much to use*

For adults aged 18 to 65, the dose is one injection of 6 mg, just under the skin, on the outside of the thigh.

If the first *Imigran* Mk II injection helps your migraine or cluster headache, but the headache/migraine comes back later, you may use another *Imigran* Mk II injection. You must wait at least one hour between using the first and second *Imigran* Mk II injection. Do not use more than 2

injections (2 x 6 mg) in any twenty-four hours.

Do not use a second injection, or any other form of *Imigran*, if the first injection has not provided any relief from your symptoms. You may take your usual headache relief medication provided it does not contain ergotamine or methysergide. If you are not sure what to do, ask your doctor or pharmacist.

If your migraine or cluster headache is not relieved by *Imigran*, you may use *Imigran* Mk II injection on another occasion to treat another attack. Provided there are no side effects, you can use *Imigran* Mk II injection to treat at least three separate migraine or cluster headache attacks before you and your doctor decide this medicine is ineffective for you.

### *How to use it*

***Imigran* Mk II injection should be injected just under the skin on the outside of the thigh using the autoinjector (see the patient instruction leaflet inside the pack).**

### *When to use it*

It is best to use your *Imigran* Mk II injection -

- a. when the migraine headache or cluster headache begins; or
- b. when other symptoms of the migraine begin, such as nausea (feeling sick), vomiting or your eyes becoming sensitive to light.

If you use *Imigran* Mk II injection later during the attack it will still work for you. Do not use your *Imigran* Mk II injection before the above symptoms occur.

---

## What do I do if I use too much? (Overdose)

---

**Immediately telephone your doctor or Poisons Information**

**Centre (telephone 131126) for advice, if you think you or anyone else may have used too many *Imigran* Mk II injections, even if there are no signs of discomfort or poisoning. You may need urgent medical attention.**

**If you are not sure what to do, contact your doctor or pharmacist.**

---

## While you are using *Imigran* Mk II injection

---

### *Things you must do*

**Tell your doctor if, for any reason, you have not taken your medicine exactly as directed.**

Otherwise, your doctor may think that it is not working and change your treatment unnecessarily.

### *Things you must not do*

**Do not give this medicine to anyone else, even if their symptoms seem similar to yours.**

### *Things to be careful of*

As with many other medicines, *Imigran* Mk II injection may cause drowsiness in some people.

**Be careful driving or operating machinery until you know how *Imigran* Mk II injection affects you.**

If you use *Imigran* Mk II Injection too often, it may make your headache worse. If this happens, your doctor may tell you to stop taking *Imigran* Mk II Injection.

---

## What are the side effects?

---

**Check with your doctor as soon as possible if you think you are experiencing any side effects or allergic reactions due to using *Imigran* Mk II injection, even if the problem is not listed below.**

Like other medicines, *Imigran* Mk II injection can cause some side

effects. If they occur, they are most likely to be minor and temporary. However, some may be serious and need medical attention.

A common side-effect with *Imigran* Mk II injection is temporary pain at the site of injection.

**Tell your doctor if you experience any of the following after using *Imigran* Mk II injection:**

- pain, tingling, heat or flushing in any part of the body.
- feeling of sleepiness, dizziness or tiredness.
- nausea (feeling sick) or vomiting.
- a change in blood pressure.
- feeling of faintness.
- problems with your eyesight.
- pain in the lower tummy and bloody diarrhoea (ischaemic colitis).
- shaking or tremors,
- uncontrolled movements,
- shortness of breath.

**Tell your doctor immediately, or seek urgent medical attention, and do not use any more *Imigran* Mk II injection if you:**

- feel heaviness, pressure or tightness in any part of the body including the chest or throat.
- feel irregular heart beats.
- have a fit or convulsion.
- have wheezing, swelling of the lips/mouth, difficulty in breathing, hay fever, lumpy rash ("hives") or fainting. These could be a symptom of an allergic reaction.
- have persistent purple discolouration and/or pain in the fingers, toes, ears, nose or jaw in response to cold.

These side effects are likely to be serious. Stop using *Imigran* Mk II injection and seek medical attention straight away.

This is not a complete list of all possible side effects. Others may occur in some people and there may be some side effects not yet known.

**Do not be alarmed by this list of possible side effects. You may not experience any of them.**

---

## How do I store *Imigran* Mk II injection?

---

**Keep your autoinjector and *Imigran* Mk II injections where children cannot reach them, such as in a locked cupboard.**

**Keep *Imigran* Mk II injection in a cool, dry place where it stays below 30°C and protect from light.**

**Do not leave in a car, on a window sill or in a bathroom.**

**Keep *Imigran* Mk II injection in its pack until time to use.**

**Return any unused or expired medicine to your pharmacist.**

---

## Product description

---

### *What Imigran Mk II injection looks like*

*Imigran* Mk II injection is a pre-filled 0.5 mL volume syringe containing 6 mg sumatriptan (as succinate), in solution.

Each syringe is designed to be loaded into and used with the autoinjector device.

*Imigran* Mk II injection is packed in a box containing two syringe cartridges. Each syringe contains one dose of *Imigran*.

*Imigran* Mk II injection is also available as an Autoinjector Kit.

### *Ingredients*

*Imigran* Mk II injection contains the active ingredient sumatriptan succinate.

*Imigran* Mk II injection also contains sodium chloride and water for injection.

Latex Allergy - The needle shield of the pre-filled syringe may contain

dry natural latex rubber that has the potential to cause allergic reactions in latex sensitive individuals.

### *Supplier*

Your *Imigran* Mk II injection is supplied by:

Aspen Pharmacare Australia Pty Ltd  
34-36 Chandos Street  
St Leonards, NSW 2065  
Australia.

### *Where to go for further information*

Pharmaceutical companies are not in a position to give people an individual diagnosis or medical advice. Your doctor or pharmacist is the best person to give you advice on the treatment of your condition. You may also be able to find general information about your disease and its treatment from books, for example in public libraries.

This leaflet was updated in December 2017.

The information provided applies only to: *Imigran*® Mk II injection.

®*Imigran* is a registered trade mark of Aspen Global Incorporated.

*Imigran*® Mk II Injection  
Sumatriptan (as succinate) 6 mg/0.5 mL: AUST R 44465.

*Imigran*® Mk II Injection  
Autoinjector Kit Sumatriptan (as succinate) 6 mg/0.5 mL: AUST R 44466.

© 2003 Aspen Global Incorporated.

Version 3.0