

ALVESCO®

ciclesonide

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about ALVESCO.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

The information in this leaflet was last updated on the date listed on the final page. More recent information on the medicine may be available.

You should ensure that you speak to your pharmacist or doctor to obtain the most up to date information on this medicine. You can also download the most up to date leaflet from

<http://www.astrazeneca.com.au/CMI>

Those updates may contain important information about the medicine and its use of which you should be aware.

All medicines have risks and benefits. Your doctor has weighed the risks of you using this medicine against the benefits he/she expects it will have for you.

If you have any concerns about using this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine. You may need to read it again.

What ALVESCO is used for

ALVESCO is inhaled into the lungs for the treatment of asthma. Asthma is a disease where the lining of the lungs becomes inflamed (red and swollen), making it difficult to breathe.

ALVESCO contains a medicine called ciclesonide, which belongs to a family of medicines called corticosteroids.

ALVESCO metered dose inhalers deliver a measured amount of ciclesonide, which acts directly on your air passages to reduce inflammation and help keep the air passages open. In doing so, this medication helps to improve your condition and to prevent asthma attacks from occurring. It is a 'preventer' and needs to be taken regularly.

ALVESCO will NOT give you immediate relief of wheezing or breathlessness during sudden, severe or continuous asthma attacks. In this situation you should use a quick acting 'reliever' inhaler, which contains a different medicine, a so-called bronchodilator.

Your doctor may prescribe ALVESCO to replace corticosteroid tablets, which will mean that you need to take both medicines for a short period of time. It is important to follow your doctor's instructions in using your medicines

Your doctor may have prescribed ALVESCO for another reason. Ask your doctor or pharmacist if you have any questions about why ALVESCO has been prescribed for you.

ALVESCO is not addictive.

This medicine is available only with a doctor's prescription.

Before you use ALVESCO

When you must not use it

Do not use ALVESCO if you have an allergy to:

- any medicines containing ciclesonide
- any of the ingredients listed at the end of this leaflet

Some symptoms of an allergic reaction include skin rash, itching, shortness of breath or swelling of the face, lips or tongue, which may cause difficulty in swallowing or breathing.

Do not use ALVESCO to treat an acute asthma attack.

Do not give ALVESCO to a child under the age of 6 years.

Do not use ALVESCO after the expiry date (EXP) printed on the pack or if the packaging is damaged or shows signs of tampering.

If it has expired or is damaged return it to your pharmacist for disposal.

If you are not sure whether you should start using ALVESCO, contact your doctor.

Before you start to use it

Tell your doctor or pharmacist if you have any allergies to:

- any other medicines
- any other substances, such as foods, preservatives or dyes

Tell your doctor if you are being treated for any fungal, bacterial or viral infections.

Some treatments may interact with ALVESCO.

Tell your doctor if you are pregnant, intend to become pregnant.

Like most medicines, ALVESCO is not recommended for use during pregnancy. If there is a need to consider ALVESCO during your pregnancy, your doctor will discuss the risks and benefits of using ALVESCO during pregnancy.

Tell your doctor if you are breastfeeding or plan to breastfeed.

Your doctor will discuss the possible risks and benefits of using ALVESCO during breastfeeding.

Tell your doctor if you currently have an infection.

If you take ALVESCO while you have an infection, the medication may hide some of the signs of an infection. This may make you think, mistakenly, that you are better or that it is not serious.

Tell your doctor if you have or have had tuberculosis (TB).

Tell your doctor if you have or are currently suffering from liver disease.

If you have not told your doctor about any of the above, tell them before you start using ALVESCO.

Using other medicines

Tell your doctor if you are using any other medicines, including any that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and ALVESCO may interfere with each other. These include:

- medicines used to treat fungal or viral infections, such as ketoconazole, itraconazole, ritonavir, nelfinavir

These medicines may affect how well ALVESCO works.

How to use ALVESCO

If you are not sure how to use an inhaler, ask your doctor or pharmacist.

Children should only use their inhaler on medical advice and with the help of an adult.

The correct way to use your ALVESCO inhaler is explained on the instructions for use leaflet that comes inside the ALVESCO package. If you do not use your ALVESCO inhaler correctly your lungs might get less medicine than needed. So please read these instructions carefully before you start.

To make sure your ALVESCO inhaler delivers the correct dose you should clean it regularly as outlined in the instruction leaflet.

The doctor may have advised that you use a spacer device with your inhaler. This is also recommended for children 6-11 years of age.

If so, the AeroChamber Plus is considered a suitable device for use with ALVESCO.

When to use it

Use ALVESCO at the same time every day.

How much to use

Your doctor will have spoken to you about how much of your medicine you need to take and this will depend on your individual need.

Your doctor will normally start you on 80 micrograms once daily, if you are not currently using a 'preventer' medication. Otherwise your starting dose will depend on your current medication and how well your asthma is controlled.

There are 2 strengths of ALVESCO and your doctor will have chosen the one which best suits your asthma.

The number of puffs you are required to take daily will depend on the strength of ALVESCO you are using. Your doctor may adjust your dose so that you are taking the lowest dose of ALVESCO that will control your symptoms.

When your doctor is changing you from a corticosteroid tablet to ALVESCO, you will be asked to take

both medicines for a period of time. Your dose of corticosteroid tablets will gradually be reduced to the lowest possible level. It is important you follow your doctor's instructions.

If you forget to use it

If you forget to use your ALVESCO inhaler, skip the dose you missed and use your next dose when you are meant to.

Do not use a double dose to make up for the dose that you missed.

This may increase the chance of you getting an unwanted side effect.

If you are not sure what to do, ask your doctor or pharmacist.

If you have trouble remembering to use your medicine, ask your pharmacist for some hints.

How long to use it

ALVESCO helps control your asthma. Therefore you must use ALVESCO every day. Continue using the medicine for as long as your doctor tells you.

Do not stop using it unless your doctor tells you to - even if you feel better.

If you use too much (overdose)

Do not use more than the recommended dose unless your doctor tells you to.

The risk of effects from taking too many puffs of ALVESCO is low.

If you are worried that you or anyone else may have used too much ALVESCO, telephone your doctor, or the Poisons Information Centre (telephone Australia 13 11 26).

While you are using ALVESCO

Things you must do

If you have an Asthma Action Plan that you have agreed with your doctor, follow it closely at all times.

Tell any other doctors, dentists and pharmacists who are treating you that you are using ALVESCO.

Tell your doctor or pharmacist that you are using ALVESCO if you are about to be started on any new medicines.

If you plan to have surgery (that needs a general anaesthetic), tell your doctor or dentist that you are using ALVESCO.

If you become pregnant while you are using ALVESCO, tell your doctor or pharmacist.

Keep all of your doctor's appointments so that your progress can be checked.

Things you must not do

Do not take any other medicines for your breathing problems without checking with your doctor.

Do not give ALVESCO to anyone else, even if they have the same condition as you.

Do not use ALVESCO to treat any other complaints unless your doctor or pharmacist tells you to.

Do not stop using ALVESCO, or lower the dosage, without checking with your doctor or pharmacist.

Things to be careful of

If ALVESCO does not have its usual effect

If your asthma gets worse tell your doctor. Your doctor may tell you to take extra puffs of ALVESCO or add another medicine (such as a 'reliever puffer') to your treatment.

If your breathing suddenly becomes more difficult just after you have used ALVESCO, you will need to use a 'reliever puffer' and tell your doctor immediately.

Tell your doctor as soon as possible if:

- your 'reliever puffer' does not help your breathing as much as usual
- the effect of your 'reliever puffer' does not last as long as usual

- you need more puffs of your 'reliever puffer' than usual to get relief.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are using ALVESCO.

All medicines can have unwanted side effects. Sometimes they are serious, but most of the time they are not. You may need medical attention if you get some of the side-effects.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- nausea
- bad taste in your mouth
- mouth dryness
- irritation in your mouth or throat
- skin rash
- high blood pressure
- increased difficulty in breathing immediately following the use of ALVESCO

If you experience any of the following, tell your doctor immediately or go to Accident and Emergency at your nearest hospital:

- swelling of the lips, mouth, tongue or throat
- difficulty breathing
- severe rash or welts on your skin

You may need urgent medical attention or hospitalisation.

Very rarely, inhaled corticosteroids may make you feel anxious, depressed, irritable, or have disturbed sleep.

It is possible that some people, particularly those taking higher doses of ALVESCO for a long time may rarely suffer from the following side effects:

- rounded face
- loss of bone density

- eye problems (cataract, glaucoma)

Children and adolescents who are receiving treatment for a long period of time may have their height checked regularly by their doctor. If your growth rate is slowed your doctor may adjust your treatment.

Your adrenal glands may be affected by taking high doses of corticosteroids over a long period of time or if you change from or use high doses of oral corticosteroids. Your doctor may need to check how the adrenal glands are working.

Your doctor may also tell you to take additional oral corticosteroids during periods when your body is stressed such as during trauma, surgery and infection.

Other side effects not listed above may occur in some patients. Tell your doctor or pharmacist if you notice anything else that is making you feel unwell.

Do not be alarmed by this list of possible side effects.

You may not experience any of them.

After using ALVESCO

Cleaning

Clean and dry your inhaler each time you use it.

It is important to keep your inhaler clean and dry. It may not work as well if it gets dirty. Follow the instructions at the end of this leaflet for cleaning your inhaler.

Storage

Keep your inhaler in a cool, dry place where the temperature stays below 25°C.

Do not store ALVESCO or any other medicine in the bathroom, near a sink.

Do not leave it in the car on hot days or on a window sills.

Heat and dampness can destroy some medicines.

Keep it where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop using ALVESCO, or it has passed its expiry date, ask your pharmacist what to do with any that are left over.

Do not puncture or incinerate even when empty as the canister may explode.

Product description

What it looks like

The ALVESCO inhaler comprises an aluminium canister inside a plastic inhaler unit. The inhaler unit has a mouthpiece and a plastic cover.

ALVESCO comes in two different strengths 80 micrograms and 160 micrograms.

ALVESCO 80 inhaler is available in 120 puffs.

ALVESCO 160 inhaler is available in 60 and 120 puffs.

Ingredients

ALVESCO contains ciclesonide as the active ingredient.

It also contains:

- norflurane (a non-CFC propellant which does not deplete ozone from the atmosphere)
- ethanol

ALVESCO does not contain gluten, sucrose, lactose, tartrazine or any other azo dyes.

Supplier

AstraZeneca Pty Ltd
ABN 54 009 682 311
66 Talavera Road
Macquarie Park NSW 2113
Ph: 1800 805 342

ALVESCO® is a registered trademark of the AstraZeneca group of companies.

This leaflet was prepared in August 2016.

Australian Registration Number(s)

ALVESCO 80 micrograms:
AUST R 93724

ALVESCO 160 micrograms:
AUST R 93725

Instructions for Use

ALVESCO metered dose inhaler

Before using ALVESCO, please read these instructions carefully and follow them closely.

This leaflet does not contain all the available information. If you have any questions about ALVESCO, please refer to the Consumer Medicine Information on this leaflet. If this is a new inhaler or if you have not used your inhaler for one week or more, it must be tested before use by removing the mouthpiece cover and pressing down on the canister inside the inhaler. Release 3 puffs into the air, away from you.

You do not need to shake your inhaler before taking your medicine.

The medicine is already in a very fine solution, mixed to ensure you receive the correct dose with each puff.

Follow these instructions carefully and use the pictures to help you.

1. Remove the mouthpiece cover and check inside and outside to make sure that the mouthpiece is clean and dry.



2. Hold the inhaler upright with your forefinger on the top of the canister and your thumb on the base, below the mouthpiece.
3. Breathe out as far as is comfortable.



4. Place the mouthpiece in your mouth and close your lips firmly around it.
5. Just after starting to breathe in through your mouth, press down on the top of the inhaler to release a puff while still breathing in slowly and deeply.



6. Hold your breath, take the inhaler from your mouth and remove your finger from the top of the inhaler. Continue holding your breath for approximately ten seconds or as long as is comfortable.
7. Breathe out slowly through your mouth.
8. If you have been instructed to take another puff, wait about half a minute and repeat steps 3 to 7.
9. After use, always replace the mouthpiece cover to keep out dust. Replace firmly and snap into position.



It is important that you do not rush steps 3 to 7

A correct technique will ensure the right amount of ALVESCO is getting into your lungs every time you use your inhaler.

Care and Cleaning instructions

The mouthpiece of your inhaler should be cleaned weekly with a dry tissue or cloth.

DO NOT WASH OR PUT ANY PART OF THE ALVESCO INHALER IN WATER.

You should practice in front of the mirror for the first couple of times

until you are confident that you are operating your inhaler properly. Make sure that none of your medicine is escaping from the top of inhaler or sides of your mouth.

If you find it difficult to use the inhaler with one hand, try using both hands. Put your two forefingers on top of the inhaler and both thumbs on the base below the mouthpiece. Tell your doctor, nurse or pharmacist if you are having difficulties.

When the canister is completely empty you will not feel or hear any of the propellant being discharged.

Use of inhaler with a spacer device

If your inhaler technique is poor, your doctor may recommend the use of a spacer device in combination with your inhaler. Different brands of spacers may affect the amount of medicine delivered to lungs so it is important that you be monitored for any loss of asthma control.

Before use, the spacer should be removed from the inhaler and washed in warm water and detergent and allowed to dry without rinsing or drying with a cloth. Wash the spacer at least monthly after use.

When using a spacer, you should inhale immediately after you press down on the top of the inhaler to release a puff.

Why would your Inhaler have stopped working?

Your inhaler may not work if:

1. the mouthpiece is clogged and dirty
2. the inhaler is empty (check by shaking the canister)
3. the canister is wrongly positioned

If you have any questions about your ALVESCO medication, please refer to the Consumer Medicine Information.

Doc ID-003376760 V1.0