

# Canesten® Clotrimazole Thrush Treatment Once Cream

Clotrimazole

## Consumer Medicine Information

### What is in this leaflet

This leaflet answers some common questions about Canesten Thrush Treatment.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your pharmacist has weighed the risks of you using Canesten Thrush Treatment against the benefits they expect it will have for you.

**If you have any concerns about using this treatment, ask your doctor or pharmacist.**

**Keep this leaflet with the medicine.**

You may need to read it again.

### What Canesten Thrush Treatment is used for

Canesten Thrush Treatment is used to treat a fungal infection in the vagina and vulval area (thrush).

This medicine belongs to a group of medicines called azole antifungals.

#### What is vaginal thrush?

Candida albicans is a yeast fungus that can multiply rapidly and cause damage to the vagina when other micro-organisms aren't present in a normal balance.

Some situations which can increase susceptibility to thrush are:

- Diabetes
- Obesity
- Illness
- Fatigue
- Hormonal fluctuations (pregnancy, menstrual cycle, oral contraceptive pill, HRT)
- Prolonged use of antibiotics or steroids

Thrush may cause the following symptoms:

- Vaginal itching
- Vaginal soreness
- A white odourless discharge from the vagina (like cottage cheese)
- Pain during intercourse

The symptoms below are not usually signs of thrush:

- Abnormal or irregular vaginal bleeding
- Blood stained discharge
- Vulva or vaginal sores, ulcers or blisters
- Lower abdominal pain
- Pain when passing urine
- Fever or chills
- Foul smelling and/or unusual coloured discharge

**See your doctor if you experience any of the above symptoms.**

**Ask your doctor or pharmacist if you have any questions about whether this medicine is suitable for you.**

This medicine is available without a doctor's prescription, only through your pharmacist.

### Before you use Canesten Thrush Treatment

#### When you must not use it

**Do not use Canesten Thrush Treatment if you have an allergy to:**

- any medicine containing clotrimazole
- any of the ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include swelling of the parts of the body to which treatment has been applied; rash, itching or hives on the skin.

**Do not use this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.**

**If it has expired or is damaged, return it to your pharmacist for disposal.**

**If you are not sure whether you should start using this medicine, talk to your doctor or pharmacist.**

#### Before you start to use it

**See your doctor if:**

- this is your first infection
- you are under 18 years of age
- you have had 3 or more infections in the last 6 months
- you have allergies to any other topical treatments

**Tell your doctor if you are pregnant.**

You should see your doctor if you suspect you have thrush in the first three months of your pregnancy. Your doctor can discuss with you the risks and benefits involved with treatment.

Your doctor may also advise you use a pessary during pregnancy, instead of a cream, as this can be inserted without an applicator.

**Tell your doctor if you are breast-feeding and want to use this treatment.**

It is not known whether the active ingredient, clotrimazole, is excreted in breast milk. Your doctor will be able to discuss with you the risks or benefits involved with treatment.

**Tell your doctor if you are diabetic.**

Thrush in diabetics can mean that your blood sugar levels are not well controlled. Your doctor will be able to check this for you and advise treatment.

**If you have not told your doctor about any of the above, tell him/her before you start using Canesten Thrush Treatment.**

#### Taking other medicines

There are no reports of Canesten Thrush Treatment interfering with other medicines.

### How to use Canesten Thrush Treatment

**Follow all directions given to you by your doctor or pharmacist carefully.**

They may differ from the information contained in this leaflet.

**If you do not understand the instructions on the box, ask your doctor or pharmacist for help.**

#### How to use it

**Wash your hands thoroughly before handling the cream or using the applicator.**

To avoid the possibility of re-infection, strict personal hygiene must be observed.


The cream should preferably be inserted in the evening, just before bed time, with the use of an applicator. Using it before you go to sleep will have the best effect.

#### Loading the applicator

1. Wash your hands thoroughly.
2. Pull out the plunger of the disposable applicator until it stops.
3. Open the Canesten cream tube. Attach the applicator to the tube and hold it firmly pressed against the tube.
4. Squeeze the cream tube until the applicator is full.
5. Remove the tube of cream and replace the cap.

#### Inserting the applicator

6. Carefully insert the end of the applicator containing the cream as gently and deeply into the vagina as possible. This is best achieved when lying on your back with your knees bent.
7. Hold the applicator in position and slowly push the plunger with the forefinger until it stops. This places the cream in the vagina.
8. Remove the applicator from the vagina and dispose of the applicator.



#### How long to use it

**Use the cream once.**

This treatment is a full dose in one application. The 5 g tube contains enough cream for the one dose.

**If you are not sure what to do, ask your doctor or pharmacist.**

---

## While you are using Canesten Thrush Treatment

---

### *Things you must do*

**Discontinue treatment and tell your doctor or pharmacist if you develop a local reaction to this product.**

**See your doctor if the symptoms are not gone within 4 days.**

### *Things you must not do*

**Do not take Canesten Thrush Treatment to treat any other complaints unless your doctor tells you to.**

**Do not give your treatment to anyone else, even if you suspect they have the same condition as you.**

**Do not stop using this medicine until you have completed the course.**

If you stop taking it suddenly, your condition may worsen.

### *Things to be careful of*

**Preferably the treatment should be timed to avoid the menstrual period.**

This will allow the treatment to be most effective. Ask your doctor if you have questions about this.

**Canesten Thrush Treatment may reduce the effectiveness and safety of latex products, such as condoms and diaphragms.**

This effect is temporary and occurs only during treatment.

### *Things that may help to resolve Thrush*

Some self help measures suggested below may help your condition. Your doctor or pharmacist can give you more information about these measures.

- **Blot with toilet paper after urinating (instead of wiping back to front).**
- **Change underwear, night attire, towels and linen daily.**

### **Avoid where possible:**

- perfumed toilet paper
- fragrant soaps and bath oils
- wearing tight or synthetic underwear
- douches or vaginal deodorants

---

## Side effects

---

**Tell your doctor or pharmacist if you experience discomfort while you are using Canesten Thrush Treatment.**

This medicine helps most people with thrush but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

**Do not be alarmed by the following side effect. You may not experience it.**

**Ask your doctor or pharmacist to answer any questions you may have.**

**Tell your doctor or pharmacist if you notice any of the following and they worry you:**

- skin reaction to the cream (stinging, redness, blistering, peeling, swelling, itching)

Any of the above may be signs of an allergic reaction. Your doctor or pharmacist may recommend you discontinue treatment if you experience these effects.

---

## After using Canesten Thrush Treatment

---

### *Storage*

**Keep your cream in the tube until it is time to use it.**

If you take the cream out of the tube it may not keep well.

**Keep your cream in a cool dry place where the temperature stays below 25°C.**

**Keep the treatment where children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

### *Expiry*

The expiry date is marked on the end panel of the carton and is also printed on the crimp at the end of the tube. The date refers to the month and year of expiration eg 0805, refers to the eighth month of 2005. The product can be used until the end of this month.

### *Disposal*

**If your doctor tells you to stop using this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.**

**Discard the medicine out of the reach of children.**

---

## Product description

---

### *What it looks like*

Canesten Thrush Treatment Once Cream is a white opaque cream presented in a screw -cap tube containing 5 g of cream.

The package also contains 1 disposable plastic applicator.

### *Ingredients*

Canesten Thrush Treatment

Once Cream contains 100 mg clotrimazole as the active ingredient in each gram of cream. It also contains:

- Sorbitan monostearate
- Polysorbate 60
- Cetyl palmitate
- Cetostearyl alcohol
- Octyldodecanol
- Benzyl alcohol
- Purified water

This medicine does not contain sucrose, gluten, tartrazine or any other azo dyes.

### *Manufacturer*

Canesten® Thrush Treatment Once Cream is made in Spain.

Canesten® Thrush Treatment is distributed in Australia by:

Bayer Australia Limited  
(ABN 22 000 138 714)  
Consumer Care Business Group  
875 Pacific Highway  
PYMBLE NSW 2073

Toll Free: 1800 023 884

Canesten® Thrush Treatment is distributed in New Zealand by:

Bayer New Zealand Limited  
Consumer Care Business Group  
Argus Place  
Glenfield  
AUCKLAND NZ

Toll Free: 0800 847 874

® = Registered Trademark

This leaflet was prepared in August 2005.  
AUST R number 68570