

## Aldiq cream 5%

Imiquimod

### Consumer Medicine Information

#### What is in this leaflet

This leaflet answers some common questions about Aldiq. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist. If you need any more information about Aldiq, please ask your doctor or pharmacist. All medicines have risks and benefits. Your doctor has weighed the risk of you using Aldiq against the benefits it can have for you. If you have any concerns about using this medicine, ask your doctor or pharmacist.

**Keep this leaflet with the medicine as you may want to read it again.**

#### What Aldiq is used for

Aldiq cream contains the active ingredient imiquimod. Imiquimod is an immune response modifier. It activates immune cells in the body. The immune cells then proceed to kill and remove the virus-infected or cancer cells. Although the exact way that Aldiq works is unknown, it is believed to be due to its effects on the immune system.

Aldiq is used to treat solar keratosis (SK) on the face and scalp, superficial basal cell carcinoma (sBCC), and genital/perianal warts. Solar keratoses are thickened, scaly patches of skin caused by too much sun exposure. Solar keratosis is also known as actinic keratosis.

Basal cell carcinoma is a type of skin cancer. External genital/perianal warts (condyloma acuminata) are warts that appear on the surface of the penis or vulva (external female sexual organ) and around the anus.

Aldiq is not recommended for use in patients under the age of 18 years as there have been no studies of its effects in this age group.

**If you are not sure why you are using this medicine ask your doctor.**

Your doctor may prescribe Aldiq for another purpose.

#### Before using it

##### When you must not use Aldiq

##### Do not use Aldiq:

- if you are allergic to imiquimod or to any of the ingredients listed under "Product Description"
- on skin where there are open sores or wounds. Do not start using Aldiq until after the area has healed.
- on warts inside the vagina or inside the anus or inside the urethra (where you pass urine). The use of Aldiq cream on these areas has not been studied.
- on areas that are sunburnt
- if the packaging is torn, or shows signs of tampering.
- after the "expiry" date shown on the box or sachet.

##### Before you start to use Aldiq

##### Tell your doctor if:

- you are pregnant or intend to become pregnant.
- you are breastfeeding or intend to breastfeed.

- you are or you think you are HIV positive.
- you are taking any other medicines
- you have previously used Aldiq cream or other similar preparations to treat your condition.
- you have problems with your immune system
- you have an abnormal blood count
- you currently have, or in the past have had, any other medical conditions.

#### Using Aldiq

##### Dosage

Aldiq is used to treat several skin conditions. Your doctor will tell you where to apply Aldiq cream, how often and for how long to apply it for your condition. Aldiq should be applied just before bedtime and left on the skin for 6-10 hours. Sufficient cream should be applied to cover the treatment area.

**Talk to your doctor if your condition reappears after treatment.**

##### Solar Keratosis

If your doctor has prescribed Aldiq for the treatment of SK, the usual dosage is once a day, at bedtime, three times a week. For three times a week application, Aldiq can be applied, for example, on Monday, Wednesday and Friday, or Tuesday, Thursday and Saturday.

- Your doctor may tell you to continue applying Aldiq cream for 4 weeks, followed by a period of 4 weeks without any treatment. Your doctor will then check your skin condition. If any SK lesions remain, the treatment should be repeated for another four weeks.
- Alternatively, your doctor may want you to continue applying Aldiq cream for up to 16 weeks. Each treatment should continue for no more than 16 weeks at a time.

##### Superficial Basal Cell Carcinoma

If your doctor has prescribed Aldiq for the treatment of sBCC, it should be applied to the affected area once a day at bedtime for five consecutive days per week (Monday to Friday), or as recommended by your doctor. Treatment should continue for 6 weeks unless your doctor tells you otherwise.

##### External Genital or Perianal Warts

If your doctor has prescribed Aldiq for the treatment of external genital or perianal warts, it is to be applied once a day, at bedtime, three times a week or as recommended by your doctor.

For three times a week application, Aldiq can be applied on Monday, Wednesday and Friday; or Tuesday, Thursday and Saturday. Treatment should continue until the warts are completely gone. Aldiq should not be used for more than 16 weeks at a time. It usually takes 8-10 weeks for your warts to disappear but warts may clear as early as 4 weeks. If your warts reappear, talk to your doctor. If your skin reacts badly to Aldiq, your doctor may recommend that you stop treatment for a few days. It is not necessary to make up the doses that you missed or to prolong the treatment period.

#### How to use it

1. Before applying Aldiq, wash your hands and treatment area with mild soap and water and allow it to dry thoroughly. Open a sachet of Aldiq and squeeze some cream onto your fingertip.
2. Apply a thin layer of Aldiq cream onto the treatment area and rub it gently into the skin until the cream vanishes. For sBCC lesions enough cream should be applied to cover the lesion and about 1cm of surrounding skin.

One sachet contains enough cream to cover a treatment area of 20 square centimetres (approximately 3 square inches).

3. If using the sachet after application of the cream, carefully dispose of the unused Aldiq cream in the sachet where children cannot reach it. Wash your hands with mild soap and water.
4. Leave Aldiq cream on the affected area for 6 to 10 hours. Do not shower or bathe during this time.
5. After 6-10 hours, the area where Aldiq cream was applied should be washed with mild soap and water.

#### While you are using Aldiq

- Do not use more than the recommended amount of Aldiq cream. A thin layer that completely covers the treatment area and surrounding skin is enough.
- Do not cover the treatment area with bandages or other dressings after you have applied Aldiq cream. Cotton gauze dressings are okay to use, if needed.
- Stop using Aldiq cream if you become pregnant, and do not use it if you are breastfeeding.

#### If you are using Aldiq to treat SK:

- Aldiq cream should not be applied in or near the eyes, lips or nostrils.
- Do not use sunlamps or tanning beds, and avoid going into the sun as much as possible during treatment with Aldiq cream. Wear protective clothing if you go outside during daylight.
- New SK lesions may develop during treatment with Aldiq cream. These lesions may resolve during the treatment period. Even though the initial SK lesions may clear with treatment, new lesions may develop in the future and require further treatment. Aldiq cream is not a cure, since SK is considered to be a chronic skin condition.
- There are not enough data to support the use of Aldiq to treat SK on the hands and arms.
- Aldiq cream should not be applied to an area greater than 25 square centimetres.

#### If you are using Aldiq to treat sBCC:

- Aldiq cream should not be applied in or near the hairline, eyes, ears, nose or lips.
- Do not use sunlamps or tanning beds, and avoid going into the sun as much as possible during treatment with Aldiq cream. Wear protective clothing if you go outside during daylight.

- Visit your doctor regularly if you are treating more than one sBCC lesion at the same time.

*If you are using Aldiq to treat genital warts:*

**Genital warts are very contagious, therefore avoid sexual (genital, anal or oral) contact.**

If you do decide to have sexual relations, apply Aldiq cream after, not before, sexual activity. If you have already applied the cream, it should be washed off before sexual activity.

Aldiq cream may weaken condoms and diaphragms; therefore, the cream should be washed off before using a condom or diaphragm during sexual activity. Alternate forms of contraception should be considered.

The effect of Aldiq cream on the transmission of genital warts is not known. Uncircumcised men with warts under the foreskin should pull the foreskin back each day and wash underneath it. If daily washing under the foreskin is not carried out, tightness of the foreskin may occur. Early signs of tightness include swelling and wearing away of the skin, or difficulty in pulling back the foreskin. If these symptoms occur, stop the treatment immediately and call your doctor.

Female patients should take special care if applying Aldiq cream at the opening of the vagina because local skin reactions on the delicate moist surfaces can result in pain or swelling, and may cause difficulty in passing urine.

Do not use Aldiq for more than one course if you have problems with your immune system either due to illness or because of the medicines you are already taking. If you think this applies to you, talk to your doctor.

**Aldiq cream has been prescribed for your use only. Do not give it to anyone else even if you think their condition is the same as yours.**

*If you forget to use Aldiq*

If you forget a dose, apply the missed dose of cream as soon as you remember and then continue on the regular schedule. Do not double the dose at any time to make up the missed dose.

---

## Side effects

All medicines can have side effects and this may occur with the normal use of Aldiq. Tell your doctor as soon as possible if you do not feel well while you are using Aldiq. More common side effects:

The following effects have occurred during clinical trials with imiquimod cream. If the following occur and are severe enough to worry you, tell your doctor.

- Application site reactions including redness, wearing away of the skin, flakiness, swelling, hardening under the skin, small open sores, crust that forms during healing, small bubbles under the skin, itching, burning, pain, tenderness, irritation, rash, soreness, stinging, sensitivity, skin colour becomes lighter, bleeding, lumps on the skin, infection and pimples. Most of these skin reactions are mild to moderate, and are signs that the product is working. If your skin reacts badly or the skin reaction

becomes too uncomfortable when using Aldiq cream, wash the cream off with mild soap and water and contact your doctor. Your doctor may recommend that you stop treatment for a few days.

- Flu symptoms, tiredness, fever, headache, diarrhoea, back pain, muscle pain, and swollen glands in the neck, armpit and groin.

Some patients have experienced changes in skin colour (lighter or darker) in the area where Aldiq cream was applied. These changes may be permanent in some cases.

**Please tell your doctor if you experience other side effects or you do not feel well while you are using Aldiq.**

Other side effects not listed above may also occur in some people.

*If you use too much Aldiq*

Using too much Aldiq cream could cause severe skin reactions. If too much cream is applied, simply wash away the extra cream with mild soap and water. When any skin reaction has settled, you may then continue with your treatment.

**If you or someone else accidentally swallows Aldiq cream, contact your doctor or Poisons Information Centre (in Australia call 131126; in New Zealand call toll free 0800 POISON or 0800 764 766) or go to the Accident and Emergency department at the nearest hospital.**

If the cream is accidentally swallowed, nausea, vomiting, headache, muscle pain, and fever could occur.

---

## After using it

### Storage

Aldiq cream should be kept in a cool dry place where the temperature stays below 25°C. Do not freeze.

**Do not store Aldiq cream or any other medicine in the bathroom or near a sink. Do not leave it in the car or on window sills. Keep Aldiq sachets where children cannot reach them.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

### Disposal

**If your doctor tells you to stop using Aldiq or it has passed its expiry date, ask your pharmacist what to do with any that is left over.**

---

## Product description

Aldiq cream is a white to slightly off white cream, packed in a single-use foil sachet. This is enough cream to cover a treatment area of 20 cm<sup>2</sup>. In Australia Aldiq cream is supplied in boxes of 6 and 12 sachets.

### Ingredients

Each 250mg of Aldiq cream 5% contains 12.5mg of the active ingredient, imiquimod.

It also contains isostearic acid, benzyl alcohol, cetyl alcohol, stearyl alcohol, white soft paraffin, polysorbate 60, sorbitan monostearate, glycerol, methyl hydroxybenzoate, propyl hydroxybenzoate, xanthan gum and purified water.

---

## Sponsor

Aspen Pharmacare Australia  
34-36 Chandos Street  
St. Leonards  
NSW 2065

AUST R 201801.

This leaflet was revised in September 2017.