

VELETRI®

(VEL-e-tree)

Epoprostenol Sodium (E-poe-PROST-e-nol SOE-dee-um) 500 mcg /1.5 mg Powder for Injection

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about VELETRI. It does not contain all the available information.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you using this medicine against the benefits they expect it will have for you.

If you have any concerns about using this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

What VELETRI is used for

VELETRI is used to treat some types of pulmonary arterial hypertension (PAH). PAH is characterised by high blood pressure in the blood vessel that carries blood from the heart to the lungs, and increased resistance in the blood vessels of the lung. The cause of PAH is not known however there are a number of diseases such as scleroderma that are associated with PAH. VELETRI belongs to a group of medicines called prostaglandins.

VELETRI works by widening the blood vessels in the lungs and so lowering the blood pressure in your lungs (known as a vasodilator action).

Your doctor however, may prescribe VELETRI for another purpose.

Ask your doctor if you have any questions about why it has been prescribed for you.

VELETRI is not addictive.

Before you use VELETRI

When you must not use VELETRI

Do not use VELETRI if you have ever had an allergic reaction to epoprostenol or any of the ingredients listed at the end of this leaflet.

Symptoms of an allergic reaction may be mild or severe. They usually include some or all of the following: wheezing, swelling of the lips/mouth, difficulty in breathing, hayfever, lumpy rash ("hives") or fainting.

Do not use VELETRI if you have heart disease with shortness of breath, and swelling of the feet or legs due to fluid build-up.

Do not use VELETRI after the expiry date [EXP.] printed on the pack.

If you use it after the expiry date has passed, it may not work as well.

Do not use VELETRI if the packaging is torn or shows signs of tampering.

If you're not sure whether you should be using VELETRI, talk to your doctor.

Before you start to use VELETRI

You must tell your doctor if:

- you are allergic to foods, dyes, preservatives or any other medicines.
- you are pregnant, trying to become pregnant, or breastfeeding.
- you are taking any medicine to prevent blood clots, such as heparin, warfarin, aspirin or other anti-inflammatory pain killers (NSAIDs).

- you are taking any medicines that are used to treat high blood pressure, or a group of medicines known as nitrates that are used to treat angina.
- you are taking digoxin, a medicine used to treat heart failure.
- you are taking any other medicines, including medicines you buy without a prescription from a pharmacy, supermarket or health food shop.

Some medicines may affect the way others work. Your doctor or pharmacist will be able to tell you what to do when using VELETRI with other medicines.

How to use VELETRI

VELETRI will be given as an intravenous infusion only, normally through a permanently fitted intravenous catheter (during initial treatment a 'peripheral line' may be used which is a non-permanent catheter). Before VELETRI is used, it must be dissolved only in sterile water for injection or Sodium Chloride 0.9% solution for injection.

How much to use

Initial treatment with VELETRI will be carried out in a hospital. Your doctor will start you on an infusion and slowly increase the dose (every 15 minutes) to find the most effective or largest dose you can tolerate.

During this part of the treatment you will also learn about how your body reacts to VELETRI.

Your doctor will then continue the infusion based on this dose, and may increase or decrease your infusion rate depending on your response to the treatment. All changes should be done gradually and under the direction of a doctor, except in emergency situations.

If you develop pulmonary oedema (water on the lungs) during this time, your doctor may choose not to treat you with VELETRI.

How to use VELETRI

Your VELETRI infusion will be given to you as continuous intravenous infusion only, normally through a permanently fitted intravenous catheter through a pump. There are only certain pumps which can be used. Your doctor will make sure you are using the right one.

Your doctor or nurse will have shown you how to keep your catheter clean, and the area around it clean and free from infection. They will also show you how to prepare and administer VELETRI and how to stop treatment if necessary. It is very important you follow their instructions carefully.

How long to use it

Use VELETRI for as long as your doctor advises you to. VELETRI is generally used over a prolonged period of time, possibly years. It should not be stopped suddenly. Symptoms of suddenly stopping VELETRI include dizziness, weakness and difficulty breathing.

If you use too much (overdose)

As VELETRI has vasodilatory action, overdose may lead to low blood pressure,

loss of consciousness, nausea, diarrhoea, vomiting, facial flushing, headache and fast heart beat.

In hospital, the effects of VELETRI are monitored carefully by your doctor. In the unlikely event that you receive too much, appropriate action, such as reducing the dose can be taken promptly.

If you are using VELETRI at home and you think you have used too much, immediately telephone your doctor or the nearest hospital casualty department, even if there are no signs of discomfort. You may need urgent medical attention.

Keep telephone numbers for these places handy.

While you are using VELETRI

Things you must do

Tell your doctor or pharmacist that you are using VELETRI if you are about to start on any new medicines.

Tell your doctor if you become pregnant or are trying to become pregnant.

Tell your doctor if, for any reason, you have not used your medicine exactly as prescribed.

Otherwise, your doctor may think that it was not effective and change your treatment unnecessarily.

Things you must not do

Do not stop using or change the dose without first checking with your doctor.

Do not give this medicine to anyone else, even if their symptoms seem similar to yours.

Do not use VELETRI to treat any other complaints unless your doctor says to.

Things to be careful of

As with many other medicines, VELETRI may cause dizziness/ drowsiness/tiredness in some people.

Be careful driving or operating machinery until you know how VELETRI affects you. If you are affected, do not drive or operate machinery.

Side Effects

Check with your doctor as soon as possible if you have any problems while taking VELETRI, even if you do not think the problems are connected with the medicine or are not listed in this leaflet.

Like other medicines, VELETRI can cause side effects in some people.

If you think you are having an allergic reaction to VELETRI while you are receiving it, TELL YOUR DOCTOR IMMEDIATELY.

Symptoms usually include some or all of the following:

- wheezing
- swelling of the lips/mouth
- difficulty in breathing
- hay fever
- lumpy rash ("hives")
- fainting

Tell your doctor at once if you experience any of the following while you are receiving VELETRI:

- fever
- fatigue
- chills/flu-like symptoms
- facial flushing or paleness
- fast heart beat
- shortness of breath
- diarrhoea
- feeling or being sick
- wind or tummy discomfort
- anxiety, nervousness and agitation
- dizziness, especially on standing
- headaches
- dry mouth
- rash
- decreased or increased feeling or sensitivity, especially in the skin
- tingling or numbness of the hands or feet
- chest pain and tightness
- reddening and pain at the infusion site
- sweating
- jaw, muscle and/or back pain
- bleeding: any bleeding can be serious, if this occurs you should contact your doctor

Do not be alarmed by this list of possible side effects.

You may not experience any of them.

This is not a complete list of all possible side-effects. Others may occur in some people and there may be some side-effects not yet known.

Tell your doctor if you notice anything else that is making you feel unwell, even if it is not on this list.

Ask your doctor if you don't understand anything in this list.

VELETRI may affect your blood sugar levels, heart rate or blood pressure during infusion. Your doctor will monitor these.

It is very important to keep the area around the intravenous catheter clean, otherwise infection of the skin at the site of injection may result, which can then spread into your blood (known as septicaemia).

During administration of VELETRI the intravenous catheter may become blocked. Tell your doctor or pharmacist if this happens.

Storing VELETRI

Storage

Keep this medicine where young children cannot reach it.

A locked cupboard at least one-and-a half metres above the ground is a good place to store medicines.

Keep VELETRI powder in a cool, dry place where it stays below 25°C.

Protect from light by keeping it in the carton until use.

Do not store VELETRI powder in a bathroom or near a sink.

Do not leave VELETRI powder in the car or on window sills.

Do not freeze VELETRI powder at any time.

Do not use this medicine after the expiry date which is stated on the carton after

EXP. The expiry date refers to the last day of that month.

The reconstituted solution should be immediately further diluted to the final concentration. VELETRI diluted to the final concentration in the drug delivery reservoir as directed can be stored for up to 8 days at 2° to 8°C. Do not expose this solution to direct sunlight.

Do not use this medicine if you notice any particles in the reconstituted solution.

Product description

What it looks like

VELETRI: A Sterile, lyophilised white to off-white powder in a clear glass vial with a rubber stopper and an aluminium flip-off cap. Each pack contains one vial. The 500 mcg vial has a white flip-off cap and the 1.5 mg vial has a red flip-off cap.

Pack size

Each pack contains one vial.

VELETRI - 500 mcg vial of epoprostenol

VELETRI - 1.5 mg vial of epoprostenol

Ingredients

Active ingredient:

VELETRI contains the active ingredient epoprostenol, as the sodium salt.

Inactive ingredients:

- sucrose,
- L-arginine,
- sodium hydroxide (for pH adjustment)

Supplier

VELETRI® is supplied in Australia by Actelion Pharmaceuticals Australia Pty Limited.

Suite 6, 13b Narabang Way
Belrose NSW 2085

VELETRI® is distributed in NZ by Pharmacy Retailing (NZ) Ltd t/a Healthcare Logistics

58 Richard Pearse Drive
Airport Oaks, Auckland NZ.

This leaflet was prepared in September 2016

VELETRI 500mcg - AUST R 208316

VELETRI 1.5 mg - AUST R 207547