

Ceftazidime-AFT

Ceftazidime Powder for Injection 1g and 2g

Consumer Medicine Information

What is in this leaflet?

Please read this leaflet carefully before you use Ceftazidime-AFT.

This leaflet answers some common questions about Ceftazidime-AFT. It does not contain all of the available information.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Sometimes new risks are found even when a medicine has been used for many years. Your doctor has weighed the expected benefits of you taking Ceftazidime-AFT against the risks this medicine could have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

What is Ceftazidime-AFT used for?

Ceftazidime-AFT belongs to a group of medicines called cephalosporins.

Ceftazidime-AFT is used to treat a wide range of infections caused by bacteria. These infections may affect the chest (bronchitis or pneumonia), ears, nose, throat, bladder and urinary tract, skin and soft tissue, stomach or bones.

Ceftazidime-AFT works by killing the bacteria that cause these infections.

Your doctor may have prescribed Ceftazidime-AFT for another reason.

There is no evidence that Ceftazidime-AFT injection is addictive.

Before you are given Ceftazidime-AFT

When you must not be given it:

You must not be given Ceftazidime-AFT if:

- you have ever had an allergic reaction to cephalosporins, penicillins or any of the ingredients listed toward the end of this leaflet. (See "Ingredients")
- the expiry date (EXP) printed on the pack has passed.
- the packaging is torn or shows signs of tampering

Tell your doctor if:

You must tell your doctor if:

- you are allergic to foods, dyes, preservatives or any other medicines (in particular cephalosporins, penicillins, chloramphenicol or lignocaine)
- you are taking any other medicines, including medicines you buy without a prescription
- you are taking oral contraceptives (birth control pills)
- you have had to stop taking another medicine for your infection
- you have kidney or liver problems
- you have had stomach or bowel illness (for example, colitis)
- you test your urine for sugar
- while on Ceftazidime-AFT, you need to have your blood tested for a blood transfusion

- you are breastfeeding, pregnant or trying to become pregnant.

How Ceftazidime-AFT is given

How much to give

Ceftazidime-AFT will be administered as directed under the supervision of a doctor or nurse. You will not be expected to give yourself this medication.

How it is given

Ceftazidime-AFT will be injected deep into muscle tissue or into a vein. When it is injected into a vein it may be given directly or through a 'drip'.

DO NOT try to use the injection on your own unless your doctor has told you to do so and given you complete instructions.

How long it is given for

Ceftazidime-AFT should be administered for as long as recommended by your doctor.

Do not stop taking Ceftazidime-AFT, or change the dose without first checking with your doctor.

Use in children and elderly patients

Consult your doctor before using Ceftazidime-AFT in children or elderly patients.

What do I do if I am given too much? (Overdose)

Immediately telephone your doctor or Poisons Information Centre (telephone 131126) for advice, if you think you or anyone else may have taken too much Ceftazidime-AFT, even if there are no signs of discomfort or poisoning.

If you are not sure what to do, contact your doctor, pharmacist or nearest hospital.

While you are being given Ceftazidime-AFT

Things you must do

Tell your doctor if you are allergic to cephalosporins, penicillins or other antibiotics.

Tell your doctor if you have kidney or liver problems.

Tell your doctor that you are being given Ceftazidime-AFT before taking any other prescribed medicine or performing any urine or blood tests.

Some medicines may affect the way other medicines work

Things you must not do

Do not give this medicine to anyone else, even if their symptoms seem similar to yours.

Do not use Ceftazidime-AFT to treat any other complaints unless your doctor says to.

Things to be careful of

Be careful driving or operating machinery until you know how Ceftazidime-AFT affects you.

What are the side-effects?

Check with your doctor as soon as possible if you think you are experiencing any side effects or allergic reactions due to taking

Ceftazidime-AFT, even if the problem is not listed below.

Like other medicines, Ceftazidime-AFT can cause some side-effects. If they occur, they are most likely to be minor and temporary. However, some may be serious and need medical attention.

Tell your doctor immediately if you notice any of the following after receiving Ceftazidime-AFT:

- skin trouble such as a rash, red spots, itching, skin lumps, hives, blisters or peeling
- swelling of the eyelids, face or lips
- diarrhoea (even several weeks after your Ceftazidime-AFT treatment)
- high temperature (fever)
- difficulty breathing
- bruising or bleeding
- shaking or muscle twitches
- Wheezing, swelling of the lips/mouth, difficulty in breathing, hayfever, lumpy rash (hives) or fainting. These could be symptoms of an allergic reaction.

Tell your doctor if you notice any of the following after receiving Ceftazidime-AFT:

- pain or flaking skin where you had the injection
- headache
- stomach pain
- dizziness
- numbness or tingling
- nausea (feeling of sickness) or vomiting
- bad taste in mouth
- sore mouth or tongue
- vaginal discharge, genital itching.

This is not a complete list of all possible side-effects. Others may occur in some people and there may be some side-effects not yet known.

Tell your doctor or pharmacist if you notice any side effects from your medicine which are not mentioned here.

Do not be alarmed by this list of possible side-effects. You may not experience any of them.

How do I store Ceftazidime-AFT?

Keep this medicine where children cannot reach it, such as in a locked cupboard.

Store your Ceftazidime-AFT injection below 25°C.

Do not leave in a car, on a window sill or in the bathroom.

You will find an expiry (or "use by") date on the manufacturer's label of the pack. The injection should not be used after this date.

Product description

What Ceftazidime-AFT looks like

Ceftazidime-AFT looks like a white or almost white crystalline powder. The powder is reconstituted with sterile water prior to injection. Once made up the solution should be a clear and light yellow to amber colour.

Ingredients

Ceftazidime-AFT contains the active ingredient Ceftazidime pentahydrate.

Ceftazidime-AFT also contains anhydrous sodium carbonate.

Supplier

AFT Pharmaceuticals Pty Ltd
113 Wicks Road
North Ryde
NSW 2113
Australia

Where to go for further information

Pharmaceutical companies are not in a position to give people an individual diagnosis or medical advice. Your doctor or pharmacist is the best person to give you advice on the treatment of your condition.

This leaflet was revised in September 2017.

Ceftazidime-AFT Injection 1g: AUST R
277680

Ceftazidime-AFT Injection 2g: AUST R
277679